



**EPILEPSY
FOUNDATION®**
VIRGINIA

Not another moment lost to seizures™



Virginia School Nurse Seizure Update

EPILEPSY FOUNDATION OF VIRGINIA.....	6
ACKNOWLEDGEMENTS:.....	7
INTRODUCTION.....	8
Section 1	10
Seizure and Epilepsy Basics	10
Epilepsy Etiology and Prognosis.....	11
Risk of Epilepsy in Special Populations.....	11
Classifying Seizures	12
Partial Seizures	12
<i>Simple Partial Seizures</i>	<i>13</i>
<i>Complex Partial Seizures</i>	<i>13</i>
Generalized Seizures	13
Absence Seizures	14
Atypical Absence Seizures	14
Atonic Seizures	15
Clonic Seizures	15
Myoclonic Seizures	15
Tonic Seizures	15
Tonic-Clonic Seizures	15
Epileptic Syndromes	16
Benign Rolandic Epilepsy	16
Childhood Absence Epilepsy	17
Juvenile Myoclonic Epilepsy.....	17
Lennox-Gastaut Syndrome.....	17
Temporal Lobe Epilepsy	18
Frontal Lobe Epilepsy.....	18
Progressive Myoclonic Epilepsy	19
Landau-Kleffner Syndrome	19
Reflex Epilepsy	19
Seizure-Like Phenomena:	20

Medical Conditions that Mimic Epilepsy	20
Non-epileptic Seizures	20
Intractable Epilepsy	21
Status Epilepticus:	22
Prolonged or Serial Seizures	22
Nonconvulsive Status Epilepticus	23
AND EPILEPSY BASICS	23
Epilepsy in Female Students	23
THE IMPACT OF EPILEPSY	25
Impact of Epilepsy on Learning	25
Assessing Learning and Behavioral Disorders	26
Impact of Epilepsy on the Brain.....	26
Assessing the Impact of Epilepsy: The Spectrum of Severity	27
Uncomplicated.....	28
Compromised	28
Devastated	28
Behavioral Concerns	28
The Stigma.....	29
Teenage Problems	29
Section 3	32
FIRST AID	32
Basic First Aid:	32
Care and Comfort.....	32
Care of Students with No Change in Consciousness.....	32
Care of Students with Altered Awareness.....	33
Care of Students with Seizures that Cause Loss of Consciousness	33
After a Seizure Episode.....	34
Acute Interventions and Emergency Response.....	34
Delegation of Acute Intervention Responsibilities	37
Section 4	39
TREATMENT FOR SEIZURE PREVENTION.....	39
Antiepileptic Drugs (AEDs).....	39

Initiating Drug Treatment.....	39
Antiepileptic Drugs in Use.....	39
PRN Medications for Prolonged or.....	41
Acute Repetitive Seizures	41
The Cognitive and Behavioral Effects of Antiepileptic Drugs	42
Discontinuing Treatment.....	42
Ketogenic Diet	43
Modified Atkins Diet.....	44
Vagus Nerve Stimulator (VNS)	44
Service Dogs	44
Alternative Therapies	46
Managing Stress	46
What Students Can Do for Themselves.....	46
What Teachers Can Do to Help	47
Advances in Treatment	47
Section 5	49
SEIZURE ACTION PLANS	49
Medication Documentation and Administration.....	50
Documenting Seizures and Circumstances	50
Behaviors Before a Seizure	51
Is There a Pattern?	51
What Happens to the Student During a Seizure?.....	52
What Was the Length of the Seizure?	52
Behavior After the Seizure?.....	52
Observing Responses to Treatment.....	53
Assessing Learning and Psychosocial Issues	53
Customizing an Action Plan.....	53
Educational Accommodations	53
Special Circumstances and Precautions	54
Teaching and Tailoring Interventions.....	55
Supportive Counseling and Referrals.....	55
Section 6.....	58

TRAINING TEACHERS AND OTHER	58
SCHOOL PERSONNEL	58
General Curriculum for Training Others in the School Setting	58
Training Objectives	58
Considerations in Curriculum Development	58
Dispelling Myths and Reducing Fears	59
Teaching About the Impact of Epilepsy on Learning	59
Managing Seizures in the Classroom	60
Managing Seizures Under Special Circumstances	60
<i>On the School Bus</i>	60
<i>For a Student in a Wheelchair</i>	60
<i>When Seizures Occur in Water</i>	61
<i>In Non-Classroom Areas of the School</i>	61
Section 7	63
EPILEPSY RESOURCES	63
Community Resources	63
Internet Resources	63
Fact Sheets on Epilepsy	66
Driving and Epilepsy	66
Appendices	68
Appendix A:	69
Seizure Action Planning Forms	69
Appendix B:	74
Seizure Recognition and First-Aid Charts	74
Appendix C:	79
Antiepileptic Medications Chart	79
Index	84

EPILEPSY FOUNDATION OF VIRGINIA



The Epilepsy Foundation of Virginia (EFVA) is a statewide affiliate of the Epilepsy Foundation of America (EFA). It was established in 1978 to promote awareness about epilepsy, advocate for research, find a cure and provide assistance to the 80,000-100,000 Virginia and their friends and families who live with epilepsy.

The EFVA provides free training on managing epilepsy and seizure first aid for professionals, families and colleagues. Services for Virginians include medication assistance, transportation assistance and help with epilepsy related emergencies. A network of regional directors and volunteers provides support through phone and email helplines and support group information is available. Programs are scheduled through the year in localities around the state to raise awareness and funds for services and epilepsy research. Information about epilepsy is available online or by calling the EFVA office. The EFVA also writes a newsletter that is free for subscribers.

The EFVA office is located at:

500 Ray C. Hunt Drive, Charlottesville, VA 22903

The mailing address is:

P.O. Box 800659, UVA Health Center, Charlottesville, VA 22908

Telephone: 434-924-8669

Website: <http://www.efva.org>

Five regional offices and director names, addresses and contact information can be found by calling or on the website at: <http://www.efva.org>

Suzanne Bischoff is the Executive Director and can be emailed at : srb3m@virginia.edu

ACKNOWLEDGEMENTS:

We wish to thank the GENAN FOUNDATION, THE EPILEPSY FOUNDATION AND ITS COWORKERS, SCHOOL NURSE JOANNA COLLINS AND EPILPESY NURSE MALLIE HAYNES FOR THEIR CONTRIBUTIONS. WE ALSO THANK STUDENTS COURTNEY AND INDIA, AS WELL AS THE EPILEPSY FOUNDATION OF VIRGINIA PROFESSIONAL ADVISORY BOARD FOR THEIR CONTRIBUTIONS. LAST, BUT NOT LEAST, WE WISH TO THANK THE VIRGINIA DEPARTMENT OF EMERGENCY SERVICES TERRI COY AND EPILEPSY HERO MATTHEW MAYS FOR COORDINATING THE MANUAL AND DVD.

SUZANNE BISCHOFF, PH.D.

EXECUTIVE DIRECTOR

EPILEPSY FOUNDATION OF VIRGINIA



INTRODUCTION

“The school nurse’s role is multifaceted. It means being a caregiver, advocate, fact-finder and communicator. ”

This Virginia school nurse update is largely based on the 2009 EF manual with permission. We hope this update will help you in your job.

Suzanne Bischoff
Executive Director
Epilepsy Foundation of Virginia

As parents, grandparents and healthcare professionals, we can agree that all children should be afforded every opportunity to learn and to play, exploring life both at home and at school. This is how youngsters acquire both the cognitive and social skills, in conjunction with neurodevelopmental growth, needed to function in today's increasingly complex and demanding society. As William Wordsworth noted, "The child is father of the man," so the legacy of lessons learned in childhood - both positive and negative – play a pivotal role in the quality of life each child will eventually enjoy as an adult.

While most seizures last from a few seconds to a minute or two, the child who experiences them in school is often compromised on several levels. Cognitive function is often affected. But it is not the only complicating factor. These students must often cope with added challenges related to the adverse effects of drug therapy. There can be emotional and behavioral consequences as well. This multiplicity of issues adds to the complexity of managing the student's needs and safeguarding the student both physically and emotionally in the school setting.

For those working with students with epilepsy, there are many challenges. These include coping with the unpredictability of seizures and learning how to respond to a seizure.


Perhaps most importantly, however, is the school nurse's ability to help create a positive atmosphere among teachers, staff and fellow students with respect to the student with seizures. This helps to minimize stigmatizing behaviors and to support a positive self-image for the student. The school nurse can facilitate this understanding and enhance awareness and sensitivity by educating everyone in the school setting about the nature and management of seizures. By example, the school nurse can reinforce the need for compassion, patience and understanding for the student who has seizures.

While the school nurse is dearly important in influencing positive societal attitudes that we hope will prevail in the future, his/her intervention is especially crucial for the here and now. All of the school nurse's activities assist the student with epilepsy in achieving success in school life and in social relationships. The school nurse can help educators recognize the causes of behavior problems, if these occur, and can help diminish the educational and social impacts often associated with childhood epilepsy. The school nurse can also assist in early diagnosis by alerting parents and healthcare providers about possible seizure-related behaviors and offer guidance on obtaining information and services that will benefit students and their families.

Section 1

SEIZURE AND EPILEPSY BASICS

“Epilepsy is a chronic disorder, the hallmark of which is recurrent unprovoked seizures.”



Section 1

Seizure and Epilepsy Basics

Epilepsy is clinically defined as two or more unprovoked seizures occurring more than 24 hours apart in someone older than one month of age. It is the most common treatable serious neurological disorder in children and is the third most common of all serious neurological disorders in childhood, with mental retardation and cerebral palsy being the most common (Shinnar and Pel lock, 2002). It affects from four to ten school-aged children per 1,000.

A seizure is the clinical manifestation of an abnormal and excessive synchronized discharge of cerebral neurons. It is sudden and transient and can include a wide variety of motor, psychic and sensory phenomena. There may or may not be an alteration in consciousness or awareness. The location of the initial electrical disruption, the extent of its spread within the brain and its duration determine the clinical manifestations of the seizure and its impact on the student.

A seizure can include jerking movements of the limbs, falling down, string, automatisms (purposeless repetitive actions) such as lip smacking or hand wringing or even strange feelings in one's stomach. Seizures are usually unprovoked and unpredictable. The child can't control his or her behavior during a seizure.

About 10 percent of all individuals will experience a seizure during their lifetimes. These seizures may be related to acute medical, neurological or neurosurgical conditions and do not recur once the underlying cause is resolved.

Some children have a single seizure, the cause of which is never identified. These acute episodes are *not* considered to be epilepsy.

Students often outgrow epilepsy and are seizure free as adults. The majority of seizures are not emergencies, but a part of daily life. In some cases, adults may experience seizures, but most are not emergencies. Some students have more than one type of seizure. Many students have complete or near complete control when they take their medication as prescribed.

TABLE 1: Epilepsy Statistics

- Epilepsy affects up to 1 percent of all children age 16 or younger.
- As many as 325,000 school-aged children (ages 5 to 14) have epilepsy (Shinnar and Pellock, 2002).
- Up to 45,000 new cases of epilepsy are diagnosed annually in children (200,000 total new cases are diagnosed annually in the United States).
- Incidence is highest in those under age 2 and over age 65.
- In children, the median age of onset is between 5 and 6 years old.
- Incidence is greater in African-American and socially-disadvantaged populations.
- In more than 70 percent of those with epilepsy, no specific cause for the disorder can be identified.

Epilepsy Etiology and Prognosis

Epilepsy is a brain disorder characterized by the tendency to have recurrent unprovoked seizures. It may or may not be associated with damage in brain structure. In some cases, usually the more benign forms, genetics seem to play a role. Sometimes epilepsy is related to a congenital malformation in brain structure or an inborn error in brain metabolism. About 30 percent of cases of epilepsy can be attributed to trauma, infection, postnatal vascular lesions or VNS degenerative conditions. About 20 percent of cases are associated with neurological conditions presumed to be present from birth, such as mental retardation and/or cerebral palsy (see Table 3). About 70 percent of all new cases in children are considered idiopathic (assumed to be genetic) or cryptogenic (see Table 2) and have no apparent cause (Hauser, 2001)

In general, those who experience idiopathic seizures have a better prognosis in terms of both seizure control and eventual remission than do those with epilepsy symptomatic of underlying brain damage or disease. About 70 percent of those children who are otherwise normal will outgrow epilepsy and be seizure-free as adults (Shinnar and Pellock, 2002).

In childhood, the majority of seizures are partial in nature, with specific epileptic syndromes, such as benign rolandic epilepsy and childhood absence epilepsy, accounting for a large proportion of cases (Shinnar and Pellock, 2002).

Some forms of childhood epilepsy are defined as benign. They are self-limiting and often remit by adolescence. The child's development and intellect *will* most likely be normal. In other cases, although development and intellect do not appear to be affected, seizures persist despite treatment. Others are more serious and can be associated with developmental delay or cognitive impairment and persistent seizures. The prognosis depends upon a variety of factors as listed in Tables 4 and 5.

TABLE 2: Causes of Epilepsy

For approximately 70 percent of children diagnosed with epilepsy the cause is determined to be either.

- Cryptogenic (of unknown cause)
- Idiopathic (presumed to be genetic)

For the remaining 30 percent the causes are generally the same as those for symptomatic seizures.

Risk of Epilepsy in Special Populations

The basic underlying risk of developing epilepsy is about 1 percent. Individuals in certain populations are, however, at greater risk. Epilepsy can be expected to develop in:

- 10 to 15 percent of children with mental retardation
- 13 percent of people with Cerebral Palsy (McDermott, 2005)
- 8.7 percent of children of mothers with epilepsy
- Up to 50 percent of children with both mental retardation and cerebral palsy
- 2.4 percent of children of fathers with epilepsy
- 33 percent of children who have had a single seizure (Hauser, 1990)

Classifying Seizures

Seizures are classified by their clinical symptoms, supplemented by electroencephalographic (EEG) data. Accurate classification of seizures has been greatly improved by the simultaneous use of EEG and videotape recordings. However, since most children experience seizures in a non-medical setting, the most important basis for classification is still an eyewitness account.

The International League Against Epilepsy (ILAE) Commission on Classification and Terminology first proposed a classification system for epileptic seizures in 1981. Today, it is the most widely used classification system. It divides seizures into two major groups - **partial** and **generalized** (see Table 6).

TABLE 3: Causes of Symptomatic Seizures

- Brain trauma (auto accidents are #1)
- Brain lesions (e.g. tumors, tubers)
- Brain injury at birth
- Congenital malformations
- Infections of the brain (e.g. meningitis, encephalitis)
- High fever
- Metabolic abnormalities (e.g. hypoglycemia, hypocalcemia, hyponatremia)
- Poisoning

Partial Seizures

Partial seizures are the most common type of seizure experienced by children with epilepsy. Virtually any movement, sensory or emotional symptom can occur as part of a partial seizure, including complex visual, auditory or olfactory hallucinations.

In partial seizures the electrical disturbance is limited to a specific area of one cerebral hemisphere (side of the brain). Partial seizures are subdivided into simple partial seizures (in which consciousness is retained) and complex partial seizures (in which consciousness is impaired or lost). Partial seizures can spread to cause a generalized seizure, in which case the classification category is partial seizures secondarily generalized.

TABLE 4: Factors Predictive of Good Prognosis

- Normal neurological exam
- Normal intellectual functioning
- Absence of brain lesions
- Late onset of seizures (child is older than 3 or 4 years)
- Normal EEG
- Infrequent seizures
- Early response to a single medication
- No tonic and/or atonic seizures
- No status epilepticus

Simple Partial Seizures

Simple partial seizures result from abnormal neuronal activity in specific, limited areas of the brain that affect movement, sensation or emotion. Partial seizures can progress through several stages that reflect spread of the abnormal neuronal firing to different areas of the brain. For example, a seizure that begins in a motor area can cause focal twitching of the hand or face

This explains why simple partial seizures do not affect consciousness. The child remains aware of the environment, remembers the seizure experience, but may be limited in how he or she can interact while it is in progress. Speech, for example, may be arrested.

A simple partial seizure can also be manifested as tingling in the hand or face, visual distortions or hallucinations, a feeling of sudden fear, a sense of unexpected familiarity (*déjà vu*) or unfamiliarity (*jamais vu*), a breeze on the skin, a sudden rising feeling in the stomach or sudden unexplained pain. Simple partial seizures of this type are also referred to as *auras*, internal experiences that may herald more extensive seizures to come and serve as warning to the affected individual.

Simple partial seizures may be confused with acting out, mystical experiences, mental illness or psychosomatic illness.

Complex Partial Seizures

Complex partial seizures, like simple partial seizures, begin in one area of the brain, but usually involve larger portions of the brain and are associated with an alteration in consciousness. In practical terms, this means that the person experiencing the seizure cannot interact normally with his environment, has impaired responsiveness and usually does not remember what happened during the episode.

Automatisms are a common characteristic of complex partial seizures. These are repetitive, complex movements that are purposeless, undirected and inappropriate. Common examples of automatisms include lip smacking, repetitious swallowing or chewing, fidgeting with fingers or hands and repeated clumsy movements of a preceding motor act. Automatisms are likely to be stereotyped-that is, the same type of movement occurs each time the student has a complex partial seizure.

In some cases, complex partial seizures generate activities that cause social distress - disrobing, running, screaming, flailing and exhibiting excessive fear. These phenomena, while far less common than the facial and fumbling automatisms described above, nevertheless create major social difficulties for children who experience them and for their families.

Complex partial seizures may be confused with drunkenness, aggression or drug abuse.

TABLE 5: Factors Predictive of Seizure Persistence

- Abnormal neurological exam
- Brain abnormality
- Abnormal EEG
- Multiple seizure types
- Frequent seizures

Generalized seizures affect both cerebral hemispheres (sides of the brain) from the beginning of the seizure. They produce loss of consciousness, either briefly or for a longer period of time.

Absence Seizures

Absence seizures typically last just several seconds. There is no warning before the seizure and the child typically resumes whatever activity he or she was involved in before the seizure occurred. For example, if a student is reading he or she will stop. The child will usually blink his eyes a few times and the eyes may even roll up a bit. They may also have associated automatisms. When the seizure is over the student will continue to read. These seizures are extremely brief in duration and are typically not associated with the risk of injury. These seizures can occur many times a day or in clusters, both of which interfere with a child's ability to learn and function.

Absence seizures may be confused with daydreaming, ADD or inattentiveness.

Atypical Absence Seizures

Atypical absence seizures are different from typical absence seizures. They last longer, onset is not as abrupt and incomplete loss of awareness and muscle tone changes are more severe. There may be associated head drops (atonic seizures) or myoclonic seizures (see below). Atypical absence seizures usually occur as part of a mixed seizure disorder.

TABLE 6: Classification of Epileptic Seizures

I. Generalized Seizures (Convulsive or nonconvulsive)

- a. Absence
- b. Atonic/akinetic
- c. Myoclonic
- d. Tonic, clonic and tonic-clonic

II. Partial (Focal Seizures)

- a. Simple partial seizures (consciousness is not impaired)
 - i. With motor symptoms
 - ii. With sensory symptoms (including vision changes, bodily sensations, dizziness and imagined sounds, smells and tastes)
 - iii. With autonomic symptoms
 - iv. With psychic symptoms (including dysphasia, dysmnesia, hallucinatory and affective changes)
 - v. Compound (i.e., mixed) forms
- b. Complex partial seizures (consciousness is impaired)
 - i. Simple partial seizures followed by loss of consciousness .
 - ii. With impairment of consciousness at the outset
 - iii. With automatisms

III. Unclassified

IV. Prolonged or repetitive seizures (status epilepticus)

Summarized from the International Classification of Epileptic Syndromes, 1981

Atonic Seizures

Atonic seizures produce an abrupt loss of muscle tone. Other names for this type of seizure include drop attacks, astatic or akinetic seizures. They produce head drops, loss of posture or sudden collapse. Because they occur abruptly without warning, and because those who experience them fall with force, atonic seizures can result in injuries to the head and face. Children with this type of seizure sometimes wear protective headgear to avoid injury. These seizures tend to be resistant to drug therapy.

Clonic Seizures

Clonic seizures are always symptomatic of underlying brain disease. They involve rhythmic jerking movements of the arms and legs, sometimes on both sides of the body. They are uncommon and occur most frequently in neonates, infants and younger children.

Myoclonic Seizures

Myoclonic seizures are characterized by rapid, brief muscle contractions that usually occur at the same time on both sides of the body. Occasionally, they involve one arm or foot. Observers usually see them as sudden jerks or clumsiness. A variant of the experience, common to many individuals who do not have epilepsy, is the sudden jerk of a foot during sleep (sleep myoclonus).

Tonic Seizures

Tonic seizures are particularly common in children who have Lennox-Gastaut syndrome, but they can occur in anyone. They involve tensing or stiffening of muscles in the body, arms or legs. The limbs may extend and maintain this posture for seconds to a minute or so. Consciousness is usually preserved.

These seizures occur most often during sleep. However, because they usually involve all or most of the brain and affect both sides of the body, a child who is standing when the seizure starts will often fall.

Tonic-Clonic Seizures

These seizures are the most common and best known type of generalized seizure. They begin with stiffening of the limbs (the tonic phase), followed by jerking of the limbs and face (the clonic phase).

During the tonic phase, breathing may decrease or cease altogether, producing cyanosis (bluing) of the lips, fingernail beds and face. Breathing typically returns during the clonic (jerking) phase, but it may be irregular. The clonic phase usually lasts less than a minute. Incontinence can occur. The child may bite his/her tongue or inside of the mouth during the episode. Breathing afterward may be noisy and appear to be labored.

Contrary to popular belief, nothing should be placed in the student's mouth during a seizure. It is best to turn the student on one side to help prevent choking and keep the airway clear. Following the seizure, the student will be lethargic, possibly confused, and want to sleep. Headaches sometimes occur. Full recovery takes minutes to hours, depending on the individual.

Epileptic Syndromes

Using seizure type alone to classify the type of epilepsy often overlooks other important information about the student and the seizure episodes. Classifying into syndromes takes a number of these identifying characteristics and symptoms into account, including the following:

- Seizure type
- Typical EEG recordings
- Clinical features such as behavior during the seizure
- The expected course of the disorder
- Precipitating features
- Expected response to treatment
- Genetic factors

Epileptic syndromes can be either idiopathic or symptomatic of underlying brain damage or disease. In general, idiopathic forms have a better prognosis in terms of both seizure control and eventual remission than do symptomatic forms.

Benign Rolandic Epilepsy

TABLE 7: Benign Rolandic Epilepsy

- Onset usually before age 15
- Characterized by partial seizures resulting in temporary inability to speak
- Seizure often generalizes to tonic-clonic
- Most seizures are nocturnal
- 95 percent of children experience remission by age 14

Benign rolandic epilepsy is the most common epilepsy syndrome in children. It usually begins between ages three and thirteen in neurologically normal children. Also known as benign partial epilepsy of childhood, it accounts for more than one-third of all cases of epilepsy that begin in middle childhood. There is a family history of this syndrome in 18 percent of cases and the condition is thought to be genetically determined.

Seizure episodes begin as simple partial seizures, usually with twitching in the face, tongue, pharynx of the seizures are nocturnal, occurring within a few hours after falling asleep. Seizures are usually accompanied by gurgling sounds, profuse drooling and temporary inability to speak. Consciousness is generally preserved, unless the seizure generalizes to tonic-clonic seizure, which happens often. If the child experiences these seizures only during sleep, the decision is often made not to treat with medication.

The child's neurological and other functioning is usually normal. However the EEG readings most often show a dramatic focal spike in the centrottemporal regions of the brain. Most children are seizure-free five years after onset and 95 percent of them will have undergone permanent remission by age 14 (see Table 7).

Childhood Absence Epilepsy

Absence seizures account for 2 to 4 percent of all cases of epilepsy in children. These seizures are non-convulsive and tend to occur in clusters.

TABLE 8: Childhood Absence Seizures

- 40 percent outgrow seizures
- Probably inherited
- Half of these children also experience generalized tonic-clonic seizures

Seizure onset usually occurs between ages four and eight.

Children with this syndrome are otherwise normal. About 40 percent outgrow the seizures. The syndrome has a strong genetic component which is not well understood at this time. Despite its overall benign nature, approximately half of the children with absence epilepsy can expect to have a generalized tonic-clonic seizure during adolescence. The risk is reduced if seizures are quickly controlled with medication.

The prospects for remission of childhood absence epilepsy are more likely when the child is younger at onset, the seizures are easily controlled with medication, the child does not have generalized tonic-clonic seizures and there are no other neurological problems (see Table 8.)

TABLE 9: Juvenile Myoclonic Epilepsy

- Appears at or around puberty
- Is characterized by generalized tonic-clonic, myoclonic and absence seizures
- Seizures are primarily jerking of arms and legs

Juvenile Myoclonic Epilepsy

Juvenile myoclonic epilepsy is also called Janz's syndrome and myoclonic epilepsy of adolescence. It is characterized by myoclonic, tonic-clonic and absence seizures. These seizures often occur upon awakening.

Juvenile myoclonic epilepsy generally appears at puberty (between ages 10 and 20), but may have existed prior to that time. Seizures may be precipitated by sleep deprivation, early awakening, alcohol and drug use, stress, strong emotion, photic stimulation and menstruation. This is often a lifelong condition (see Table 9).

Lennox-Gastaut Syndrome

The Lennox-Gastaut syndrome (also known as myoclonic astatic epilepsy) is characterized by a combination of factors, usually including atypical absence seizures, tonic seizures, atonic seizures and mental retardation with a distinctive slow spike-and-wave EEG. Students with this syndrome are prone to convulsive and non-convulsive status epilepticus both requiring a medical intervention. Onset is usually between ages one and five.

Some children with this epilepsy syndrome are developmentally normal when the syndrome begins, but then lose ground, sometimes dramatically. Episodes of status epilepticus contribute to this decline. By age six most of these children have some degree of mental retardation.

Children with Lennox-Gastaut syndrome typically experience more than one type of seizure. The atonic (drop attack) seizures are potentially the most troubling because of injuries caused by repeated falls. As a result, many children with this syndrome must wear protective helmets. The *tonic* seizures are most common during sleep, including nap time, whereas generalized tonic-clonic seizures occur most often on awakening. As children with the Lennox-Gastaut syndrome grow older, the types of seizures change. Among teenagers with Lennox-Gastaut, complex partial seizures are the most-common form.

Temporal Lobe Epilepsy

The temporal lobes, one on each side of the head, just above the ears, are the sites of one of the most common forms of epilepsy. Complex partial seizures with automatisms, such as lip smacking or rubbing the hands together, are the most-common seizures in temporal lobe epilepsy.

Seventy-five percent of patients with this form of epilepsy also experience simple partial seizures which may include features such as: a mixture of thoughts, emotions and feelings that are hard to describe; sudden emergence of old memories or feelings of strangeness in familiar surroundings; hallucinations of voices, music, smells or tastes and feelings of unusual fear or joy and; gastric sensations (e.g. pain, nausea, emesis). While partial seizures dominate, approximately half the people with temporal lobe epilepsy have generalized tonic-clonic seizures as well.

The seizures characteristic of temporal lobe epilepsy often begin in the deeper parts of the temporal lobe (part of the limbic system) which control emotions and memory. Memory problems may develop over time in people with this syndrome.

Frontal Lobe Epilepsy

Partial seizures beginning in the frontal lobe may produce weakness or the inability to use certain muscles, including the muscles that make it possible to talk. Sudden thrashing movements during sleep are also characteristic of frontal lobe epilepsy, as is posturing with the head jerking to one side and the arm rising with it into a brief, frozen state. Sometimes a generalized convulsion follows the slow march of these movements.

Complex partial seizures in the frontal lobe have some distinct features in contrast to those in the temporal lobes. They usually last less than a minute, are less likely to be followed by confusion or fatigue and often occur in a series or cluster. They often occur during sleep and may occur many times during the night and disrupt the normal sleep cycle. This can result in daytime sleepiness, behavioral problems and poor school performance.

Frontal lobe epilepsy has significant social effects because the seizures it generates are more likely to involve brief episodes of screaming, bicycling movements or even movements suggestive of sexual activity

Progressive Myoclonic Epilepsy

Progressive myoclonic epilepsy is a rare, genetically based form of epilepsy marked by myoclonic and tonic-clonic seizures. People with this disorder also experience difficulties with balance, muscle rigidity and progressive loss of mental abilities. A gene associated with this disorder has recently been identified.

Landau-Kleffner Syndrome

Landau-Kleffner syndrome, also a rare disorder, causes the sudden or gradual loss of the ability to understand spoken language in addition to a seizure disorder. Speech in children with this syndrome slowly declines over time. There are always epilepsy related abnormalities on the EEG, even though some children with this syndrome do not have seizures. Parents often think that the child is developing a hearing problem or has become suddenly deaf. Hearing tests, however, show normal hearing. Children may also be initially diagnosed as autistic or developmentally delayed.

The syndrome typically begins between ages three and seven. Seizures generally occur while the child is asleep and may be quite infrequent. Simple partial and tonic-clonic seizures may occur. Treatment with standard antiepileptic drugs is not very effective, but treatment with steroids has been tried with some success.

Reflex Epilepsy

Reflex epilepsy is the name given to seizures which are triggered by individual sensitivity to sensory stimulation in the environment. The most-common form is photosensitive epilepsy - seizures caused by exposure to intense or fluctuating levels of light.

Seizures in photosensitive people may be triggered by exposure to television screen flicker or rolling images, certain video games, computer monitors, alternating patterns of different colors or the intense strobe lights in fire alarms. Also, some people have reported having seizures triggered by natural light, such as sunlight flickering through trees, light glancing off water or through the slats of Venetian blinds. The condition usually begins in childhood and may be outgrown by adulthood.

The reflex response is usually absence seizures but may generalize to myoclonic seizures or generalized tonic-clonic seizures. Wearing polarized sunglasses with blue lenses has been cited as good protection against photosensitive reflex seizures.

While flashing or flickering light is the most common trigger for reflex epilepsy, rare triggers include certain sounds, music, tone of voice, reading, immersion in hot water or even eating.

*About 70% of these children
who are otherwise normal
will outgrow epilepsy and be
seizure-free as adults."*

Seizure-Like Phenomena: Medical Conditions that Mimic Epilepsy

Differential diagnosis of epilepsy is sometimes confounded by a variety of medical conditions that produce sudden symptoms resembling epileptic seizures. Some of these episodes can be caused by other medical conditions that mimic epilepsy. These include:

- Migraine and transient ischemic attacks that produce visual disturbances
- Hyperventilation, which produces tingling, lightheadedness and occasionally loss of consciousness
- Syncope (see Table 10)
- Breath-holding spells that lead to cyanosis (bluing) of the lips and jerking (usually in children under age five)
- Tics that produce jerking
- Sleepwalking and other sleep disturbances that mimic seizure symptoms
- Episodic dyscontrol syndrome (rage attacks) that are erroneously identified as epileptic seizures
- Narcolepsy, characterized by intermittent, uncontrollable episodes of falling asleep during the daytime which can produce cataplexy (short-lived intermittent muscle weakness) and sleep paralysis (paralysis while falling asleep or waking)
- Certain paroxysmal movement disorder with chorea or dystonia

TABLE 10: DIFFERENTIATING SYNCOPE/FAINTING FROM EPILEPTIC SEIZURES

CLINICAL DATA	SYNCOPE/FAINTING	SEIZURES
Precipitating factors	<i>Almost always</i>	<i>Occasional</i>
Warning	<i>Lightheaded, dizzy, queasy</i>	<i>May have aura</i>
Fall	<i>Often slumps to the floor</i>	<i>May be violent</i>
Motor component	<i>Rare</i>	<i>Frequent</i>
Incontinence	<i>Rare</i>	<i>Occasional</i>
Self-injury	<i>Rare</i>	<i>Common</i>
Appearance	<i>Pale, clammy</i>	<i>Rubor, cyanosis, diaphoresis</i>
Pulse	<i>Decreased, irregular</i>	<i>Usually increased</i>
Degree of postictal	<i>Mild</i>	<i>Often marked</i>
Family history	<i>Often positive for syncope</i>	<i>Often positive for seizures</i>
Interictal EEG	<i>Usually normal</i>	<i>Often abnormal</i>

Non-epileptic Seizures

Non-epileptic seizures or psychogenic seizures, also called pseudoseizures, are non-epileptic, episodic, paroxysmal events not related to abnormal electrical activity in the brain. Considered to be of psychological rather than physical origin, they offer a major challenge to diagnose and treat.

While these episodes do resemble true epileptic seizures, they are different from them in several important ways:

- Students have repeatedly normal EEG readings during seizure-like events.
- There is a lack of any response to therapeutic levels of antiepileptic drugs
- The seizures may be marked by violent thrashing of all four limbs, with typically asynchronous and arrhythmic movement, without losing consciousness. It is unusual in epilepsy for someone to remain

conscious in the presence of sustained motor activity of the arms and legs.

Psychogenic seizures are best dealt with on an individual basis. Once identified through video-EEG monitoring, the problem can be addressed through individual and family counseling and psychotherapy. Occasionally treatment may include psychiatric medication. Sometimes the seizures abate when the person is informed that they are psychological in nature.

TABLE 11: Differentiation Epileptic Seizures from Non-Epileptic Seizures

CLINICAL DATA	NON-EPILEPTIC SEIZURES	GENERALIZED TONIC-CLONIC SEIZURES	COMPLEX PARTIAL SEIZURES
Changes in seizure frequency with medication change	<i>Rare</i>	<i>Common</i>	<i>Common</i>
Increased seizure frequency with stress	<i>Frequent</i>	<i>Occasional</i>	<i>Occasional</i>
Combativeness	<i>Common</i>	<i>Rare</i>	<i>Rare</i>
Vulgar language	<i>Frequent</i>	<i>Rare</i>	<i>Rare</i>
Self-Injury	<i>Rare</i>	<i>Occasional</i>	<i>Rare</i>
Incontinence	<i>Rare</i>	<i>Common</i>	<i>Rare</i>
Tongue Biting	<i>Rare</i>	<i>Common</i>	<i>Rare</i>
Nocturnal occurrence	<i>Rare</i>	<i>Common</i>	<i>Occasional</i>
Stereotype of attacks	<i>Often variable</i>	<i>Little variation</i>	<i>Non identical but usually have similar patterns</i>
Postictal confusion, lethargy, sleepiness	<i>Rare</i>	<i>Always</i>	<i>Frequent</i>
EEG, Interictal	<i>Often normal</i>	<i>Frequently abnormal</i>	<i>Frequently abnormal</i>
EEG, During attack	<i>Normal</i>	<i>Always abnormal</i>	<i>Usually abnormal</i>

Intractable Epilepsy

Students with intractable epilepsy fail to respond to standard antiepileptic drug therapy. Often, they have underlying structural brain or neurological conditions, head injury or cognitive problems. These children continue to experience seizures and are usually on large amounts of multiple antiepileptic medications. They must endure not only the ongoing seizures but also the side effects of the drug therapy. Of the 326,000 children under age 15 with epilepsy, approximately 90,000 have severe seizures that cannot be adequately treated.

Students with intractable epilepsy are the greatest challenge for school nurses. They are more likely to experience seizures during school and, therefore, require more time and attention. They may require onsite emergency interventions by the nurse or a trained designee.

Although single seizures do not permanently impair intellectual or behavioral functions, multiple and uncontrolled seizures can have significant consequences. The long-term effects of numerous or recurrent seizures are the subject of studies and controversy and should never be underestimated.

Intractable seizures also significantly impact the student's quality of life, with limitations on participation in sports, schoolyard games and class outings. Students with more severe seizures may need an aide with them at all times and/or must wear a protective helmet. This can further stigmatize and shame the student and have deleterious effects on the student's self-esteem.

These students are at high risk of becoming isolated and withdrawn. The school nurse should facilitate the student's ability to participate safely in school activities and counsel the family whenever there are concerns about the student's emotional well-being.

Status Epilepticus: Prolonged or Serial Seizures

Most seizures end after a few seconds or a few minutes. If they are prolonged or occur in a series, there is an increased risk of status epilepticus. Traditionally, status epilepticus is medically defined as 30 minutes or more of uninterrupted seizure activity or more than one seizure without regaining baseline within 30 minutes. Many believe, however, that much shorter periods of continuous seizure activity may cause neuronal injury and that seizure self-termination is unlikely after as few as five minutes. More recent medical definitions consider 10 minutes, rather than 30 minutes, as the defining time. The Epilepsy Foundation advises calling for emergency assistance whenever a convulsive seizure continues for more than five minutes without signs of stopping.

Although rare, death or brain damage can occur as a result of status epilepticus. During status seizures problems can arise if there is pulmonary or cardiac arrest that is not promptly treated. More often, however, serious negative consequences occur hours or days after the onset of status as a result of prolonged stress, oxygen deprivation and systemic complications such as organ failure.

TABLE 12: Status Epilepticus Facts

- 10 percent of patients who develop epilepsy have status as their first seizure
- Of all of those who experience status, 25 percent are individuals with epilepsy
- 15 percent of persons with epilepsy experience an episode of status epilepticus

In children, the most common precipitating factors for status epilepticus are fever/infection and withdrawal from medication or noncompliance with the prescribed regimen (DeLorenzo, 1992). There may, however, be no obvious cause for the episode. Ten percent of children who later progress to epilepsy have experienced their first seizure as a status episode.

In all cases, prompt treatment is the key to preventing serious negative outcomes. The goal of treatment is to stop the seizure activity as quickly as possible and treat any underlying precipitant (DeLorenzo, 1992). Mortality in children and adults is minimized when status lasts less than one hour. After an hour, however, mortality increases slightly in children but jumps dramatically to close to 38 percent in adults.

Any type of epileptic seizure can progress to status epilepticus, but convulsive status has the greatest potential for causing long-term damage.

Treatment of convulsive status epilepticus generally includes use of such drugs as diazepam, lorazepam, phenytoin and phenobarbital administered in a planned sequence. Rectal diazepam gel may be prescribed for at-home or non-hospital use to stop bouts of prolonged seizures or clusters of acute repetitive seizures in children with a history of this type of seizure.

Nonconvulsive Status Epilepticus

Nonconvulsive status can present in various ways, including loss of speech, automatisms and alteration of consciousness. It includes continuous absence seizures and partial status epilepticus, as well as status involving simple partial seizures called *epilepsia partialis continua*.

Although nonconvulsive status epilepticus is not usually a dangerous or life-threatening condition it may have an adverse effect on a child's memory and intellectual abilities, particularly if it is not recognized early. The treatment of nonconvulsive status epilepticus, whether absence or complex partial status epilepticus, is similar to the treatment of convulsive status epilepticus.

AND EPILEPSY BASICS Epilepsy in Female Students

More than one million women and girls in the United States have epilepsy. While many of the issues they face are common to both genders, women with epilepsy must deal with unique levels of uncertainty at various points in their reproductive lives from puberty onward because of the influence of hormones associated with menstrual cycles. Epilepsy related to the menstrual cycle is called *catamenial epilepsy*. While research in this area is just beginning, it is already yielding important new information.


Additional research is needed regarding the effectiveness of contraception for women with epilepsy. All commonly used birth control methods can be used. Hormonal contraception is usually the most effective method, but it can be compromised by the type of AEDs used.

Antiepileptic drugs that induce the hepatic microsomal enzyme system that clears drugs through the liver can, in some women, lower the effectiveness of birth control pills, requiring higher doses to prevent conception. This is important for older female students to know, especially if they are sexually active.

Section 2

THE IMPACT OF EPILEPSY

*“Understanding where
a child is on the
spectrum of severity
helps determining the
extent of intervention
necessary at given
points in time”*



Section 2

THE IMPACT OF EPILEPSY

The many variables in epilepsy add to the complexity of its impact on students. Seizures may be controlled or uncontrolled. They can lead to disabling events or simply minor challenges. Medications can have mild or severe side effects. The stigma of epilepsy impacts self-concept and socialization skills. Therefore, when evaluating the consequences of epilepsy, it is important to recognize that the conjoined impact of all of these factors - seizures, medications, external and/or internalized stigma, and underlying brain abnormalities can be significant. Each variable, alone or in combination, can drastically affect the child's learning, behavior, self-concept, psychosocial development, and overall quality of life. This defines the extent to which he/she is disabled by epilepsy.

Impact of Epilepsy on Learning

Most children with epilepsy are normal and have IQs within the normal range. As a group, however, their risk for learning problems is three times greater than average. While most children with epilepsy are otherwise normal, as a group, their risk for learning problems is significantly increased. Research suggests that about half of children with epilepsy meet the criteria for a learning disability in at least one area (Fastenau et al., 2008). In this study, learning disabilities were found in all three areas: writing (38%), mathematics (20%), and reading (13%). Underachievement often occurs in language skills, reading and mathematics. The long-term adverse impact of these deficits may extend beyond education to employment, marriage and fertility (Shinnar and Pellock, 2002).

Children, who achieve seizure control relatively quickly, with few medication side effects and no cognitive impairments, generally have the best chance for average or above-average educational achievement. However, it is worth noting that children with epilepsy with average IQs may not achieve up to their potential, and attention problems are identified in many children with epilepsy.

Children with epilepsy also experience scholastic problems with memory, reaction time, concentration and visual spatial learning. Loss of school time, because of seizures or medical tests, may also affect performance, even among students who are otherwise doing well.

Other finding and observations about the impact of epilepsy on learning and behavior in children include:

- 55-65% of students with epilepsy attend regular classes. 30% require special education. 15% are "uneducable."
- Young age at onset, poor seizure control, underlying brain lesions and mixed EEG patterns are some of the predictors of the need for special education (Zelnick et al., 2001).
- When intellectual disability and seizures coexist, a higher rate of disturbed behavior is found (Sabaz, 2001).
- In a quality of life survey, intellectually normal children with refractory (uncontrolled) epilepsy had difficulty with school performance and experienced socialization problems and a reduced quality of life (Sabaz, 2001).
- Educational achievement is influenced by several factors: neurological and cognitive impairments, psychiatric problems, underlying pathology and social and cultural influences, as well as the degree of seizure control (Sperling, 2004; Hermann, 1993)

TABLE 13: Factors that May Increase the Risk of Learning, Psychosocial and Behavioral Problems

- | | |
|--------------------------|----------------------|
| • Early age of onset | • Seizures in school |
| • Frequent seizures | • Memory deficit |
| • High seizure frequency | • Slowed motor speed |

Assessing Learning and Behavioral Disorders

Students with seizures may exhibit behaviors and have learning consequences that should be assessed while determining eligibility for an Individualized Education Plan (IEP) and completing the Individualized Healthcare Plan (IHP). These may include:

- Attention deficit disorder (ADD) or attention deficit hyperactivity disorder (ADHD)
- Learning disorders
- Adjustment disorders - psychological and social adjustment to epilepsy
- Mood disorders - anxiety and depression are most common
- Mobility problems - motor deficits such as hemiparesis or spastic paresis, which are most likely caused by underlying etiology. Episodic weakness or incoordination may be due to postictal effects of seizures or side effects of medications
- Injuries - from seizures or falls, are usually minor, but may include head injuries, fractures, lacerations and bruises
- Developmental delay - seizures are often a symptom of neurological dysfunction for children with developmental delay or mental retardation

While most children with epilepsy can participate with other students in the regular classroom, some may require additional services, which are supported by federal law, known as the Individuals with Disabilities Education Act (IDEA) and Section 504 of the Rehabilitation Act.

IDEA guarantees children with disabilities a "free, appropriate public education (F APE) in the least restrictive setting" which means school districts must develop an individualized education plan (IEP) that is tailored to the individual needs of the child with the disability. All children between the ages of 3 and 21 have the right to be evaluated at least once every three years to determine if an IEP is necessary or needs to be modified. Students and parents have the right to participate in the meetings.

Section 504 guarantees reasonable accommodations to students with impairments that substantially limit one or more major life activities, such as performing manual tasks, walking or learning.

For a child with epilepsy, commonly requested services include: administering medication in the event of a prolonged seizure, specialized transportation, specialized recreation services for a child with uncontrolled seizures, and counseling to help with the social issues of epilepsy. Specialized instruction may also be necessary for accompanying learning disabilities or cognitive impairments resulting from seizures or medication.

(Epilepsy Foundation of America)

More information about the rights of children with disabilities can be found at the Virginia Department of Education's Website: http://www.doe.virginia.gov/special_ed/index.shtml

Impact of Epilepsy on the Brain

Different types of seizures can have a variable impact on the brain, lifestyle and safety. Research suggests that uncontrolled seizures can become part of a progressive disorder (Sutula, 2003), with persistent seizures resulting in increasing cognitive dysfunction and changes in the brain tissue as a part of disease progression. These problems are not seen in individuals with controlled epilepsy, which reinforces the importance of early diagnosis and effective treatment of seizure disorders.

Impact of Epilepsy on Social Development and Quality of Life

One leading expert refers to epilepsy as "unique among illnesses for the extent and degree of psychosocial disability it causes" (Sperling, 2004). Others confirm this, reporting:

- The unpredictable and incapacitating nature of seizures can cause stress and embarrassment.
- Behavior problems in children with epilepsy are frequently the result of a combination of factors, including family stress and seizure frequency (Hermann and Austin 1993). Family stress is often more significant than the number of seizures and may be negatively impacted by lack of support from the extended family, financial problems and low self-esteem. Accordingly, family issues and attitudes must be addressed as part of any treatment plan.

Students with epilepsy are more likely to experience impaired self-image, behavior problems and psychiatric co-morbidities than are students with other chronic medical problems, such as asthma. Those with frequent or uncontrolled seizures may also face social rejection by peers. The school staff needs to realize the importance of setting the standard for acceptance of children with epilepsy. If the adults react calmly and appropriately, the classmates will be more likely to react positively.

Epilepsy still carries much stigma. This often affects psychological well-being and, for many children, it influences their ability to make and keep friends and to feel confident in social settings away from their parents. The impact of epilepsy on adolescent growth and development can be dramatic. The normal transition from dependence to independence is often disrupted. Families report concerns about safety, social isolation, frequently missed school days, grades and, eventually, driving (Swartztrauber K., 2003).

Epilepsy has measurable effects on most primary areas of life including education, marriage and family and emotional adjustment. It is to be expected that the more severely affected an individual student or adult, the greater the social and psychosocial difficulties they will experience. Once seizures are under control, the psychosocial impact may outweigh the medical impact.

Assessing the Impact of Epilepsy: The Spectrum of Severity

Epilepsy affects individuals to varying degrees, creating a spectrum of disability that is broad and ranges from having a very limited effect on the individual to having a devastating impact on all aspects of life. For the majority) perhaps as high as 80 percent) of those with epilepsy, seizures can be substantially reduced or completely controlled, enabling children to live normal or close to normal lives.

For the remaining 20 percent, epilepsy is truly disabling, marked by frequent seizures, other impairments, memory and other cognitive effects and a highly compromised standard of living. Children whose seizures are not well controlled often experience social, emotional and academic problems that affect various stages of development.

These problems commonly extend into adult life, where the same issues become chronic barriers to employment and independence. For many other children and adults, the level of functioning actually falls somewhere between these extremes.

Marshall and Cupoli (1986) have identified three categories of severity among students with seizure disorders. The severity of the seizures, the effects of the medication and the neurological status of patients all interact to produce these characteristic levels of functioning.

The categories are defined as:

- **Uncomplicated**, because the seizures are controlled with medication
- **Compromised**, because of social, emotional and educational (cognitive) problems
- **Devastated**, by virtue of multiple problems

Uncomplicated

Children in this category are likely to have gained control of seizures fairly rapidly and have no, or infrequent, seizures. They have limited, if any, side effects from medication and are cognitively intact (although memory problems are commonly cited across all levels of severity). They have no additional physical or mental impairments.

While social issues based on society's reaction to epilepsy also affect children across all severity measures, epilepsy is less disabling for children with uncomplicated epilepsy. Educational/employment problems among this group may occur, but their impact is less than those with complicated epilepsy.

Compromised

Children in this category have seizures that are usually controlled and do not have serious mental or motor deficits. However, they take higher doses of medication to maintain control and are more likely to experience a greater level of social, emotional and educational/employment problems during their lives.

Devastated

Children and adults whose lives are devastated by epilepsy often have seizures as a result of brain disease or injury that also impairs learning, memory, attention and motor and emotional function. Their seizures start early in life and may never be controlled, despite the use of multiple and combined medications. They are likely to be heavily medicated and may experience some slowing or retardation. They have difficulty in maintaining family support and social relationships and require help in everyday living. They are more susceptible to bouts of continuous seizure activity (flurries) and status epilepticus. Individuals of all ages in this category have a lifelong need for services and are substantially limited in their ability to function independently.

Understanding where a child is on the spectrum of severity helps in determining the extent of intervention necessary at given points in time. It can also be a useful reference tool in determining when added assistance may be needed for a child whose epilepsy is destabilized by medication changes, puberty or other factors.

Behavioral Concerns

About 25% of children with epilepsy have significant behavior problems. Another 25% have emotional difficulties. Epilepsy, which affects the brain, can cause or worsen behavior difficulties due to:

- underlying brain damage
- the seizures themselves
- small electrical discharges between the seizures
- the effects of the seizure medications

Any of these can impair normal brain functions or cause chemical imbalances that can lead to emotional difficulties. Seizures can make children more prone to behavior problems:

- Complex partial seizures can result in hyperactivity and problems in paying attention or controlling temper.
- Seizures from the left side can cause anxiety and frustration due to problems in understanding and expressing ideas.
- Seizures from the right side can cause impulsive behavior and social difficulties.

- Seizures from the front of the brain can cause disorganization or misunderstanding of cause and effect.

The child's social and emotional development is also influenced by reactions of people around him/her. It is important that teachers and other students are aware that reactions that isolate or ostracize the child have just as much effect on the child's behavior as the seizures and medications. (Svoboda, William MD, Epilepsy.com, 26 Sept 2003)

When there are behavior problems or concerns with a student, side effects of medications should be considered first. Behavior contracts can also be implemented for behavior that the student can control. A behavior contract is an agreement between the student, teacher and parents that sets limits for behavior, rewards good choices and outlines consequences for poor choices. Samples of behavior contracts are found in Appendix A.

The Stigma

In a 2001 survey of 20,000 teens, the Epilepsy Foundation reports that:

- More than 50% had never heard or read about epilepsy.
- 71 % of teens with epilepsy have reported being bullied and met with prejudice.
- More than 50% said they would not or were not sure if they would date a person with epilepsy.
- 19% thought epilepsy was a form of mental illness.
- 52% thought that people often die from epilepsy.
- 67% admitted they would not know what to do if they saw someone have a seizure.

To reduce the stigma it is essential to incorporate epilepsy education into the school health curriculum for all students and include seizure first aid. It is also important to educate all school personnel with first aid and support strategies. The EFV A can provide free puppet shows for younger children to help them understand both the child with epilepsy and the epilepsy, itself. The EFV A can also provide free first aid training for any school group. Knowing what to expect if a seizure occurs and how to respond will make the student, the teacher and the classmates much more comfortable with the possibility.

(Epilepsy Foundation of America)

Teenage Problems

Teenagers face many challenges and teenagers with epilepsy face more, such as worrying about how others will react to seizures and never knowing when the next seizure will occur.

Common problems teens often face are cognitive problems, such as difficulty paying attention, slow thinking, finding the right words, and memory problems. Mood swings, irritability, impulsive behavior, and anxiety may also be seen. Physical symptoms, such as headaches, stomach pains, or all over pain may occur as well. One of the most important distinctions to make is how much of what occurs is due to common adolescence, and how much is due to the effect of epilepsy or medication. Also, keep in mind that common symptoms of adolescence such as mood swings or irritability may be amplified in teens with seizures.

Before a person can obtain a driver's license in the state of Virginia, they must be seizure free for six months. Persons with nocturnal seizures may drive, but are subject to review every three months. A person who has been seizure free for two years may drive without any restrictions. If the person already has a license, it will be suspended for six months from the date of the last seizure/blackout. Unfortunately, this may influence teens to deny they are having seizures in order to keep their licenses. Though admitting to having seizures can take away the license for a few months, it also makes it possible to adjust medications and become seizure free again.

Dating is a natural part of teenage social life but teens worry about how or when to tell a potential friend about their seizures. It is better for friends to be informed before witnessing a seizure without any information. Most friends will appreciate the information.

Combining alcohol and drugs with medications is never a good idea. Drinking alcohol when underage, or

using illegal drugs at any age, is not good for anyone. With epilepsy, alcohol and drugs may increase the risk of having seizures. These risks outweigh doing these things just because everyone else does.

Students with epilepsy are encouraged to play sports and take the same precautions as all other students, such as wearing helmets and not swimming alone. The doctor can help determine how to enjoy the sport safely.

Teenage girls sometimes experience more seizures in connection with hormonal changes. The doctor should be notified if seizures are worse during the time of monthly periods.

Everyone with epilepsy feels angry and sad at times. Those feelings are normal. Talking with a trusted friend or adult can help when feeling hopeless or overwhelmed.
(Epilepsy Foundation of America)

Section 3

FIRST AID



*“Basic first aid
(care and comfort)
is the responsibility
of all school personnel
who interact
with a child who
has seizures”*

Section 3

FIRST AID

The school nurse should be informed by parents or guardians if there is a history of any type of seizure for any student in his/her care. A seizure action plan should be developed that includes plans for both basic first aid and acute intervention, should these ever be needed. See Section 5 for more information about seizure action plans.

Children with epilepsy and their parents often feel safer when, not only teachers, but also classmates are aware of the possibility of seizures. With parental agreement, it is best to provide information about epilepsy to the class. The EFY A can provide free puppet shows for younger children to help them understand both the child with epilepsy and the epilepsy, itself. The EFY A can also provide free first aid training for any school group.

In the event of a seizure, classmates need to be reassured that when the child has a seizure, he/she will be fine in nearly all cases.

First aid can be broken down into two steps:

1. Basic First Aid - Care and Comfort
2. Acute Interventions and Emergency Response

Basic First Aid: Care and Comfort

Seizures can last from a few seconds to a few minutes, depending on the type of seizure. They may affect awareness, but are rarely medical emergencies and usually require only basic care and comfort. If a student has more frequent seizures than is usual for that student, or if seizures appear different than usual, the school nurse or a delegate should follow the seizure action plan and contact the parents or other designated emergency contact.

Basic first aid (care and comfort) is the responsibility of all school personnel who interact with a child who has seizures. The school district should provide a training opportunity to all relevant staff according to the student's IHP and seizure action plan so they can respond appropriately to routine seizures. Basic first-aid training can be performed by the school nurse or by staff from a local Epilepsy Foundation affiliate. Basic first aid does not need to be made complicated. Most people, with instruction, can safely provide basic first aid.

Basic first-aid measures for seizures have common elements. However, the type of assistance provided to the student who loses consciousness because of a primary or secondarily generalized seizure (see Table 15) is different than that for students who experience states of altered awareness as seen with complex partial seizures (see Table 14).

Care of Students with No Change in Consciousness

When seizures do not alter the student's awareness or consciousness, as in *simple partial seizures*, school personnel who are assisting the student should focus on ensuring the student is safe. Make sure the school nurse or trained school personnel is with the student during the event, observing and recording what transpires. After the seizure, the school nurse or trained school personnel should ensure that the student is fully oriented to person, place and time, is able to communicate his/her needs and is capable of resuming prior activities. If needed, the student should also be provided with a comfortable, safe and supervised place to rest until he/she feels able to actively and appropriately participate.

Care of Students with Altered Awareness

Absence seizures are characterized by brief lapses in awareness. A typical seizure lasts five to ten seconds. No specific first aid is required for absence seizures. However, when these occur in clusters, the student may experience a longer period of feeling spacey, out of contact or confused. In these instances the educator/school personnel should ensure that the student is fully oriented and able to resume prior activities.

Complex partial seizures last longer than absence seizures and can affect the student's awareness of what is occurring and cause confusion. Some children report that they can hear but cannot talk. Others can talk, but may not remember what they have said or their speech does not make sense.

If a complex partial seizure lasts more than five minutes beyond its usual duration for that student, or if another seizure begins before the student fully regains consciousness, call EMS or another pre-determined emergency assistance provider as noted in the student's seizure action plan .

TABLE 14: Care and Comfort for a Complex Partial Seizure

Caregivers responding to a complex partial seizure should:

- Stay calm and reassure the student that he/she is safe
- Reassure the other students
- Speak softly and calmly, keeping in mind that the student may not be able to obey verbal instructions
- Protect the student's privacy whenever possible
- Direct the student away from potentially harmful objects such as tables, chairs and doors
- Allow a wandering student to walk in a contained area
- Stay with the student until complete awareness of the environment is maintained
- After the event, reorient the student to the surroundings and provide a comfortable, safe and supervised place to rest, if needed

Caregivers should NOT:

- Restrain the student's movements
- Expect instructions to be followed
- Shout

Care of Students with Seizures that Cause Loss of Consciousness

Generalized tonic-clonic seizures, whether primary or secondarily generalized, are often called major motor seizures. Observers often believe that these seizures last longer than they actually do. For example, while the student may not be able to respond appropriately for 20 minutes or more, the actual seizure, with stiffening and/or jerking movements of the arms or legs, may last one to two minutes or less.

After a Seizure Episode

A written protocol dictating what happens after a student has a seizure should be included in the student's seizure action plan. This protocol should be a collaborative effort between the school nurse, parent(s) and the healthcare team. The majority of students will not need to leave the classroom or school after a seizure. Whether a student leaves school after a seizure is often dependent upon the final assessment of a professional school nurse. Some students will need to rest in the school nurse's office or another comfortable, safe and supervised area until they are able to return to normal activities. If the student is not able to resume scheduled activities within a reasonable period of time, as documented in the seizure action plan, the school nurse or other trained school personnel should contact the student's parent(s) or healthcare professional(s) to determine the best course of action.

TABLE 15: Care and Comfort for a Generalized Tonic-Clonic Seizure

If a student is having a generalized tonic-clonic seizure:

- Stay calm and reassure the student that he/she is safe
- Reassure the other students
- Help the student to lie down, if necessary
- Observe and time events
- Turn the student on one side
- Protect the student's head and remove eyeglasses
- Loosen any tight clothing from around the neck
- Remove potentially harmful objects from the surrounding area
- Follow the seizure action plan
- Call EMS (or other pre-determined emergency assistance) if the seizure lasts more than five minutes unless the student's typical seizure is longer as noted in the seizure action plan

Caregivers should NOT:

- Restrain the student's movements
- Put anything in the student's mouth
- Expect instructions to be followed
- Shout

Acute Interventions and Emergency Response

The school nurse should develop a seizure action plan for every student with a known seizure condition. This should include a detailed description of recommended seizure abortive therapy and appropriate emergency responses to ensure the student remains safe in the event of an acute seizure emergency, including status epilepticus.

Each school typically has a system in place for school nurses and other personnel to call for emergency medical assistance. In some school districts, 911 is available. In others, the school may have arrangements with local hospitals or healthcare systems. The school nurse should inform parents of students with seizures about emergency system options and discuss when these systems would be needed. She/he can then develop guidelines in concert with the parents and healthcare team, defining specifically what con-

stitutes an emergency for that student and how to respond appropriately. All of this information should be included in the seizure action plan.

Currently the Epilepsy Foundation recommends initiating acute interventions after five minutes of continuous seizure activity. The school nurse or other trained school personnel should also be prepared to administer (within the state law) PRN medications as prescribed and incorporated into the seizure action plan.

When is a Seizures Considered and Emergency?

Call 911 and seek emergency assistance immediately when a seizure occurs in a student who:

- Has no previous history of seizures
- Has a convulsive (generalized tonic-clonic) seizure that lasts longer than five minutes
- Experiences repeated seizures with loss of consciousness or a second seizure occurs before the student regains consciousness
- Has more seizures (of any kind) than usual or has a different kind of seizure than normally experienced as noted in student's seizure action plan
- Has an actual injury (or a suspected one) as a result of a seizure or if parents or guardians request emergency help or an evaluation
- Has diabetes, is pregnant or has another medical condition
- Does not resume normal breathing at the conclusion of the seizure

IMPORTANT: If a student is not breathing at the conclusion of the seizure, begin mouth-to mouth resuscitation immediately and call 911 or the local emergency medical personnel.

Use of Medications for Prolonged or Cluster Seizure

Physicians sometimes prescribe an antiepileptic medication for PRN use for children who have clusters of seizures (multiple seizures within a defined period of time), prolonged seizures (status epilepticus) or a tendency to have seizure emergencies. The administration of oral medications during a seizure should be avoided, but if the student is alert between seizures, a fast-acting antiepileptic drug, such as lorazepam (Ativan®) or midazolam can be given buccally or sublingually, as prescribed by the student's physician. If a student is not alert or if other routes cannot be used, diazepam rectal gel (Diastat® AcuDial™) may be used.

Medication administration for acute or prolonged seizures may be accomplished in different ways depending on a state's nurse practice act, school district policy, school policy, state laws, physician order and parent preference and should be noted in the seizure action plan. In Virginia, the law states that any school personnel who have been trained may administer medications, including diazepam. Good nursing

practice requires the school nurse to work proactively with the students, parents and health-care providers to clarify indications and instructions for the medication's use.

The student's seizure action plan should incorporate instructions for medications to be used during prolonged seizures or seizure emergencies. Seizure medication and treatment instructions should include:

- Brand and generic name of the antiepileptic medication
- Dosage
- Dosing regimen
- Special instructions for administration
- How to monitor responses
- Potential side effects
- Possible drug interactions
- What to do if a dose is missed
- What to do if side effects are observed

Vagus Nerve Stimulation and Magnet Use

Vagus nerve stimulation therapy delivers preprogrammed electrical impulses to the vagus nerve on an ongoing basis. A Vagus Nerve Stimulator (VNS), usually implanted in the upper left chest or under the arm, stimulates the vagus nerve, which in turn transmits electrical impulses to critical brain areas that are involved in seizure generation or spread.

A VNS device may be stimulated manually with the use of a magnet, typically worn on the wrist or belt. If the student has a seizure warning sign (aura), or at the onset of a seizure, the student, school nurse or trained school personnel may swipe the magnet over the VNS device to trigger an extra burst of stimulation. Many people with epilepsy report that activating the VNS device in this manner helps to abort the seizure or lessen its intensity. The magnet may be used as often as needed with at least a minute between swipes. If the magnet is held over the VNS device for six seconds it will turn it off temporarily. The student may consciously do this to avoid certain side effects, such as changes in voice or coughing during activities involving public speaking or singing or when stimulation is uncomfortable.

The most common side effect of vagus nerve stimulation is mild hoarseness and slight tingling in the neck when the signal is activated. Other reported side effects include dyspepsia, insomnia and difficulty swallowing.

A magnet should be with the student at all times. However, the student should be reminded to keep the magnet away from anything that is sensitive to magnetic fields including computers, cards with magnetic strips (e.g. credit cards) and videotapes.

Instructions for VNS magnet use should be incorporated into the seizure action plan for students with a VNS device as a first-aid intervention that can be administered by the student, school nurse or trained school personnel. Students with a VNS implant should also carry VNS identification cards.

Videos that demonstrate the use of the Vagus Nerve Stimulator are available from the manufacturer, as are magnets. Visit www.vnstherapy.com to obtain ordering information (please see page 39 for additional information on VNS therapy).

Delegation of Acute Intervention Responsibilities

Federal laws like Section 504 of the Rehabilitation Act of 1973, the 1990 Americans with Disabilities Act (ADA) and the 2004 reauthorized Individuals with Disabilities in Education Act (IDEA) mandate that reasonable accommodations be enacted in schools to ensure that a student with a disability has full access to all school activities in the least restrictive environment and that each student has educational opportunities equal to his/her nondisabled peers.

This mandate requires that both public and private schools make appropriate arrangements to accommodate a student who may need to receive PRN medications while at school or during school activities such as routine school bus rides, field trips and sporting or fine arts events. Whether this administration is accomplished by a licensed professional school nurse or a delegate depends on state laws, a state's nurse practice act, school district and school policy and finally, parent and child preference. Where state laws restrict the delegation of medications, the school nurse, individual school, school district and student's parent(s) must discuss this issue and develop a plan for how to provide the student who may require these medications with full access to all activities.

Unfortunately, even when it is determined that delegation of acute intervention responsibilities is legal and necessary, there may be no one immediately available or willing to perform this function. When this occurs, the school nurse, parents and school administrators must seek (within state law) a workable solution to ensure that an appropriately trained individual is delegated the authority and is available to perform this function.

When delegation of an acute intervention is considered, the school nurse should make certain that all involved in the delegation process (i.e. teachers, parents, school staff, classmates or other students) understand that the school nurse must:

- Put the health, safety and welfare of the student above all other considerations
- Abide by the state nurse practice act
- Abide by the school and school district policies
- Abide by applicable state and federal mandates
- Abide by state education, public health and pharmacy laws and regulations
- Determine if a nursing task can be delegated under her nursing license

If a determination is made that a school nurse can delegate a task under her nursing license, depending on state law, the nurse is then responsible for the following:

- Ensuring that the chosen delegate is appropriate
- Providing training and ongoing assessment and documentation of the competence of the delegate
- Providing ongoing assessment of the student's health

Note: In situations where a professional school nurse is not available, the training and supervision of approved delegates may fall on the student's prescribing physician.

Section 4

TREATMENT FOR SEIZURE PREVENTION

*“The goal of
treatment
for the student with
epilepsy is no seizures
and no medication
side effects.”*



Section 4

TREATMENT FOR SEIZURE PREVENTION

When treatment is indicated for the student with epilepsy, there are a variety of options, which, used alone or in combination, can help prevent seizures and minimize their impact and severity. These include:

- **Antiepileptic drugs** (AEDs) are the cornerstone of epilepsy treatment and the most common approach. (Please see Appendix C for medications used in the United States along with dosage information and the most common side effects.) PRN AEDs are essential in managing acute seizures and status epilepticus.
- **Surgery** is an option when there is a brain lesion or abnormality that can be identified as the origin of seizure activity and can be removed safely.
- The **ketogenic diet** is high in fat and low in protein, carbohydrates and fluids. It produces a ketotic state that is beneficial in reducing seizures in some individuals with epilepsy.
- The **modified Atkins diet** is similar to the ketogenic, but not quite so restrictive.
- Vagus nerve stimulation therapy sends an electric stimulus to the brain, aborting seizures.
- **Service dogs** can provide alerts and help the handler to safety in the event of a seizure.
- Alternative therapies, while not well studied, are currently under investigation by researchers.
- Strategies for **managing stress** can lower the seizure threshold.

Antiepileptic Drugs (AEDs)

75% of people with epilepsy have seizures that are controlled with medication. Before 1990, antiepileptic medication choices were limited, but since then, nine new products have been introduced. Some work well for some individuals, so it is important for the student's healthcare providers and parents to determine the best course of treatments. (See Appendix C for more information on drug therapies.)

Initiating Drug Treatment

The decision to begin drug treatment is based upon many factors, including the risk of having recurrent events. Clinical experience and research findings demonstrate that monotherapy is effective for controlling seizures in approximately 65 percent of patients. Treatment with two AEDs controls seizures in another 10 percent.

It is common practice today to start treatment with one anti epileptic drug after the child has had two unprovoked seizures. If the child does not respond to one drug, a second drug is often added. In more resistant seizures, multiple medications are added and may control seizures. Unfortunately, multiple medications can also have negative side effects. There is frequently an effect on attention, somnolence and/or cognition when medications are used in high doses or in combinations.

Antiepileptic Drugs in Use

There are many antiepileptic drugs in use today. The oldest of these are Dilantin (phenytoin), phenobarbital, Tegretol (carbamazepine), Zarontin (ethosuximide) and Depakote (valproic acid).

The newer AEDs include Felbatol (felbamate), Gabitril (tiagabine), Topamax (topiramate), Zonegran (zonisamide), Keppra (levetiracetam), Lamictal (lamotrigine), Trileptal (oxcarbazepine), Lyrica (pregabalin)

and Neurontin (gabapentin). Many of these products are available in different formulations for ease of administration. A table of AEDs with images, dosages, and side effects can be found in Appendix C.

The student's healthcare provider, along with his/her parents, will decide when treatment is necessary and which medication should be used. The prescribed treatment and medications are usually based on seizure type, frequency and severity and the ability to perform daily living tasks at home and school as appropriate to the child's age and stage of development. Titration of medication in epilepsy is routinely needed, because dosages are adjusted for weight, side effects and efficacy. When new medications are being introduced and/or current medications are being weaned, flexibility is needed. These processes can be time-consuming and can put the student at risk for seizures during the period of adjustment.

Approximately 25 percent of those with epilepsy have seizures that are not completely controlled with medications alone. It is important to make early determinations of treatment intractability and promptly refer the child to an epilepsy specialty center to minimize both the psychosocial consequences of uncontrolled epilepsy and the potential for physical sequela (Wiebe, 2004)

The primary risks related to drug therapy range from common side effects (dizziness, drowsiness or nausea) to toxicity from medication, which can be idiosyncratic, dose-related or chronic. Conversely, allowing a child to have seizures also has risks. These include physical injury associated with falls and psychosocial issues associated with restrictions placed on the child because of epilepsy, the social stigma of seizures and the child's ongoing concerns about having another seizure.

Administering AEDs at School

The administration of all prescriptive drugs should be authorized in writing by a licensed prescriber. Written authorization should include:

- student name,
- licensed prescriber's name, phone number and signature
- date the prescription was written
- name of the medication, dosage and time of day to be administered
- anticipated length of treatment
- diagnosis or reason for the medication, unless confidential
- possible serious reactions to medication
- possible serious reactions to not receiving medication
- special handling instructions
- Parental consent must be obtained before the drug can be administered. Written authorization should include:
 - student name
 - parent name, phone number and signature
 - statement and date of consent
 - allergies
 - name of medication
 - reason for medication
 - duration of treatment

The medication must be in the original container before it can be administered.

It is recommended that the school nurse or designated school personnel ensure:

- the medication is given correctly and documented
- appropriate authorization and consent forms are completed prior to giving the medication
- the medication is properly labeled and stored in a secure place

Each time medication is administered a record should be kept documenting:

- who administered it
- to whom it was given
- the manner in which it was given
- the effect of the medication
- any side effects

Any changes in dosage or type of medication should be accompanied by new authorization forms, parental consent forms and a newly labeled container from the pharmacy.

TABLE 16: Drug Reaction Warning Signs

These are not common, but serious.

- | | |
|----------------------|----------------------|
| • Rash | • Pinpoint bleeding |
| • Prolonged fever | • Weakness |
| • Severe sore throat | • Excessive fatigue |
| • Mouth ulcers | • Swollen glands |
| • Easy bruising | • Lack of appetite |
| | • Increased seizures |

PRN Medications for Prolonged or Acute Repetitive Seizures

Several fast-acting AEDs are currently prescribed for PRN use for students who have clusters of seizures (multiple seizures within a defined period of time), prolonged seizures (status epilepticus) or a tendency to have seizure emergencies. These include AEDs such as lorazepam (Ativan) and midazolam (Versed), both of which may be given buccally or sublingually, and diazepam rectal gel (Diastat AcuDial).

Diazepam rectal gel, often referred to as a seizure rescue medication, is the most commonly prescribed medication for out-of-hospital use to stop acute or cluster seizures. It is approved by the U.S. Food and Drug Administration for use by family members or other non-medical caregivers. It is supplied in pre-packaged syringes for rectal administration, with dosages tailored to the child's weight. Remember that medication administration in the school setting depends on state laws, the state's nurse practice act, school district and school policy and finally, parent and child preference. Virginia law states that any school personnel trained in administering medication may administer it.

Diazepam rectal gel is usually well tolerated. Some students may be tired after the drug is given and need to rest, while others can resume their usual activity immediately. Primary concerns about diazepam rectal gel include student privacy during administration, the ability to assess when to administer and respiratory depression. Despite concerns, respiratory depression is not a common side effect of this medication.

Videos demonstrating the use of the diazepam rectal gel are available from the manufacturer, as are rectal syringes. Visit www.diastat.com to obtain ordering information.

Midazolam treatment by intranasal routes is also indicated for the treatment of prolonged seizures in children. Although this nasal spray is not currently indicated for out-of-hospital use in the United States it shows promise as an effective seizure rescue medication in the future. In preliminary studies intranasal midazolam has been shown to be at least as effective as diazepam rectal gel in stopping prolonged and cluster seizures in children (Harbord, 2004).

The Cognitive and Behavioral Effects of Antiepileptic Drugs

In children, the side effects from AEDs can be unpredictable and must be weighed against the benefits. Some common adverse effects include double vision, sedation, dizziness and ataxia. Other effects, like depression and psychosis, are dose dependent. Carbamazepine can induce anxiety, agitation and insomnia. Valproate can produce increased appetite and weight gain. Barbiturates can produce hyperactive responses. Phenytoin and phenobarbital have been associated with alterations in personality and changes in intellectual functioning. Although the newer agents (Lamotrigine, Topiramate and Gabapentin) are less well studied than older medications, they appear to have fewer cognitive side effects.

Common AED Side Effects

Dose Related	Drug Related
Diplopia, blurred vision	Cognitive problems
Dizziness, lightheadedness	Fatigue or drowsiness
Sedation	Weight gain or loss
Slowed thinking	Cosmetic
Feeling drunk	Hyperactivity
Coordination problems	Slowed movement
Unsteady walking	Personality changes
Nausea	Mood changes

It is often difficult to determine if behavior and mood changes are related to the AEDs. The long term effects of AEDs in children are not clear but, given the developmental challenges of a normal childhood and adolescence, even mild drug effects can have an adverse impact (Drone, 2002)

Discontinuing Treatment

Antiepileptic medications prevent seizures but do not change the status of the underlying epilepsy. Research has shown that if a child is seizure-free for two years on medication, he/she is likely to remain seizure free without medication. With that in mind, children can be weaned off AEDs in a process that is usually accomplished over a period of several weeks.

During the drug withdrawal process and for some time thereafter, school personnel should be especially observant for signs of recurrent seizures. The majority of seizures recur within six months of having medication withdrawn. For children with no major risk factors for recurrence, the rate of recurrence following withdrawal of drug therapy is less than 15 percent (Shinnar and Pellock, 2002).

Surgery

About 25 percent of children have seizures that are not well controlled, even with medications. Those who have not responded to the first drug, or even a second one, have a reduced chance of obtaining control of their seizures through standard medication therapy. In these situations, the addition of third and fourth medicines is unlikely to be successful. Some of these children are candidates for surgery.

Several different surgical procedures have been shown to control seizures. The more common include focal resection, temporal lobectomy, lesionectomy and corpus callosotomy. Surgery typically offers relatively high post-surgical cure rates. Some studies suggest that post-surgical seizure-free rates are as high as 75 percent one year after surgery.

Evaluation prior to a surgical procedure is rigorous and generally includes inpatient video/EEG monitoring for five to ten days. The goal is to locate the area of the brain from which seizures are emanating as well as areas affected by seizures. Besides noninvasive and invasive EEGs, the workup often includes diagnostics such as magnetic resonance imaging (MRI), positron emission tomography (PET) and/or single photon emission computed tomography (SPECT). Neuropsychological testing may also be conducted.

Side effects can take several weeks or months to completely disappear. Typical side effects include headaches, dizziness, unsteadiness, jaw aches, swelling or bruising of head and face, blurred vision and sensation changes, especially numbness around the operation site. Depression or mood changes may also occur, but are generally temporary. Headaches often worsen with coughing, sneezing or straining. Jaw aches occur because the muscles controlling the jaw may need to be cut. Blurred vision can last for several weeks.

A child will usually go home after seven days in the hospital and will likely miss several weeks or months of school. Therefore, it is advisable to discuss post-surgical academic needs with a student and their parents prior to the surgery and develop a plan for re-entering the school environment.

Ketogenic Diet

The ketogenic diet was first introduced as a treatment for intractable seizures in the late 1920s. This diet, which is high in fat and low in protein, carbohydrates and fluids, produces a ketotic state in the child. It appears to be most effective for myoclonic seizures which involve brief, shock-like jerks of a muscle or a group of muscles. It may also be helpful for other seizure types, such as tonic-clonic and complex partial seizures. The precise mechanism of action is not completely understood.

The physician, nutritionist and nurse must collaborate to supervise elements of this dietary therapy. The child is usually hospitalized when the diet is initiated, because two to three days of fasting are needed to produce the initial ketotic state. Afterward, the diet must be strictly followed. Even a minute amount of sugar can disrupt the ketosis and jeopardize the child's seizure-free state.

Parents need to provide prepared meals and snacks for the child, who is not allowed to eat anything else. Each item provides nutrients mathematically calculated to keep the child in a state of ketosis. The diet also includes fluid restrictions. Supplemental vitamins and minerals are given in sugar-free form.

Not surprisingly, compliance is the main problem that school nurses see with students on this regimen. If an older student is doing well on the diet, the school nurse can often promote compliance by emphasizing its benefits. Younger students need to be observed carefully to prevent them from eating and drinking another student's food. Teachers and other school personnel and volunteers need education on the importance of maintaining the student's dietary restrictions.

As with other forms of treatment, the ketogenic diet does have side effects. Renal calculi and cardiomyopathy can occur. Blood abnormalities may occur but are not common.

Modified Atkins Diet

The Modified Atkins Diet is a modification of the Ketogenic Diet and was first formally studied at Johns Hopkins in 2002. Research at Johns Hopkins is still in progress, but studies so far show that 2/3 of the participants show a 50% reduction in seizures after six months. This diet is a low-carbohydrate, high-fat alternative to the ketogenic diet. There are no fluid, calorie or protein restrictions and fats are strongly encouraged. Foods are not weighed and measured, but carbohydrates are counted. This diet is easier to follow than the ketogenic diet inside and outside the home and can include the family. Side effects can include weight loss, increased cholesterol and increased risk of kidney stones. It is recommended that the physician supervise the diet with blood and urine monitoring every three months. If the diet is not helpful in controlling seizures, it should be stopped.

(Johns Hopkins Epilepsy Center)

Vagus Nerve Stimulator (VNS)

Vagus nerve stimulation therapy is usually reserved for adjunctive treatment in refractory epilepsy. This approach to managing seizures consists of implanting a generator subcutaneously in the left upper chest or under the arm. A wire conducts the electric impulse from the generator to the left vagus nerve in the neck. When the generator sends an electric signal through the wire, the vagus nerve sends this stimulation to the brain. Although the exact mechanism of action is not completely understood, these intermittent electrical signals interrupt epileptiform activity for some people.

The generator is programmed to automatically deliver intermittent stimulation to the vagus nerve. The frequency and amount of stimulation can be controlled with a programming device. Initially, the child is started on a low level of current that is increased gradually. Periodic reprogramming of the generator is generally needed, based upon seizure control and patient response to treatment. A VNS device may also be stimulated manually with the use of a magnet, typically worn on the wrist or belt. Many people with epilepsy report that activating the VNS device in this manner helps to abort the seizure or lessens its intensity. (See page 33 for more information on VNS magnet use as a seizure first-aid intervention.)

Service Dogs

Virginia state law affords each student the right to be accompanied by a service dog in a public school. This right, however, is weighed against the rights of other students at the school.

The 2008 amendments to the Virginians with Disabilities Act provide the right of the student to be accompanied by a service dog, making IEP or 504 determinations unnecessary. When requested for educational reasons, the inclusion of the service animal in the school is certainly

an appropriate matter for the student's IEP or 504 plan. When requested for noneducational reasons, the service animal's presence in the school is determined by the school division's administrators based on the school division's policy and procedures.

Any animal, pet or service animal is personal property and cannot be brought onto school property without prior approval. However, if a student arrives at school with an animal without prior approval, the animal will remain with the student until the parent or guardian arrives and removes the animal from school property. The animal will not be taken away from the student by school personnel.

The ADA regulations are clear that a service animal must be under the control of its handler. The school may request removal of the service animal if the animal is out of control or the handler does not take effective action to control it or the animal is not housebroken.

The ADA regulations state that a school division may not require proof of service animal certification or licensing, nor shall it require proof that the animal has been certified, trained or licensed as a service animal. Neither may the school make inquiries about a service animal when it is readily apparent that an animal is trained to do work or perform tasks for an individual with a disability.

The ADA regulations clarify that animals that are used purely for emotional support and not trained to perform tasks to mitigate the effects of a disability, are not service animals. Individuals with mental disabilities who use service animals that are trained to perform specific tasks, such as preventing impulsive or destructive behaviors, are protected.

The ADA regulations state that a "trained service dog" can be called a: hearing dog, guide dog, assistance dog, seizure alert dog, mobility dog, psychiatric service dog, or autism service dog. A "trained service dog" is not called a: skilled companion animal, therapy dog, social dog, facility dog, agility dog, police dog, search and rescue dog, helping dog or support dog.

School administration should also consider the factors related to service animals in the school:

- A rest place for the service animal
- fire drill or emergency evacuation plan
- pre-treating the school on a regular basis to reduce animal-borne allergens
- training for any adults who may handle the service animal
- a school-wide educational program to educate others on how to behave around the service animal

(Code of Virginia, 51.5-44)

See "Service Animal Access Request Checklist", Appendix A

Alternative Therapies

Alternative therapies for treating epilepsy have not been validated by rigorous scientific studies and their usefulness has not been established. It is important to question parents about the use of any alternative therapies, since these approaches might interfere with the effects of medical treatment.

Some complementary therapies that have been used for seizure control include acupuncture, aromatherapy, biofeedback and stress management techniques. Research regarding the effectiveness of these approaches is underway.

Managing Stress

Some common triggers for seizures include lack of sleep, flashing lights, certain foods, illness or fevers. There is no documentation showing a relationship between stress and seizures but patients often report that during periods of increased stress, they experience a greater likelihood for seizures. Stress can also affect sleep patterns and increase the risk of illness, which can be triggers for seizures. Because stress can lower the seizure threshold in so many ways, it is important to practice some basic strategies for relaxing and managing stress. The goal of relaxation techniques is to achieve a state of relaxation and reduced stress.

Physical exercise such as aerobics, cardio or weight training, and yoga can help release stress and decrease the likelihood of a seizure. A technique called Progressive Relaxation, involves doing simple motions such as curling toes or balling fists for short intervals. This helps to relax one part of the body at a time until it is completely relaxed. Whatever the workout is, individuals should always be aware of the possibility of having a seizure during a workout and take appropriate precautions.

Some herbs are believed to help with controlling seizures. However, there is not enough research to fully support this. Herbs are not regulated by the FDA and individuals should first consult their doctor if they are thinking of trying herbal remedies.

Support groups or counseling can help reduce stress, both physical and emotional. If the brain is feeling overworked, often talking to someone can help to manage stress.

Relaxation therapy to help manage stress includes taking a walk, playing music and doing things that are enjoyed like a hobby or recreational activity. Many people enjoy things such as playing sports, taking classes, reading, or writing.

What Students Can Do for Themselves

It is important for students with epilepsy to take as much control as possible. These are some ways to take control medically, cognitively and socially:

- Do not go to class if you experience warning signs of a seizure
- Inform your teacher of your seizures and be sure to explain if you miss a class because of one.
- Keep up to date with class work. Falling behind can cause higher stress levels.
- Ask for tutoring help if needed.
- Take medication every day at the same time.
- Get plenty of sleep, stay hydrated and eat well.
- Practice managing stress.

What Teachers Can Do to Help

Academic difficulties associated with epilepsy include memory problems, drowsiness, fatigue, visual problems and absenteeism. These are some ways teachers can help:

- Allow the student to tape the class or compare notes with the teacher.
- Have preferential seating for the student.
- Accommodate the student's request for a special help session.
- Be informed about the reasons for absences.
- Have open communication with the student and parents.
-

Advances in Treatment

Clinical testing of new medications continues and many of the more recently approved medications are only now being used often enough in children to provide insight into their pharmacological properties and potential. In addition, technological advances in radiology and imaging are helping to better define the anatomy and physiology of the brain.

Knowledge gained from this research will promote more precise diagnoses and create greater potential for surgical solutions for certain types of epilepsy. Research will also help provide clues to the pathogenesis of epilepsies, which in turn will aid in developing new AEDS.

Section 5 **SEIZURE ACTION PLANS**

“To ensure the student’s needs are met, a seizure action plan must be student-centered and identify strategies to overcome barriers within school settings.”



Section 5

SEIZURE ACTION PLANS

A seizure action plan is a tool used by school nurses and other school personnel to organize, communicate and tailor the healthcare and educational needs of the student with seizures. The plan helps ensure a safe environment for students, while maximizing opportunities for learning and development. In addition to documenting seizure type and treatment, this plan should document basic and emergency first-aid procedures and list specific interventions and strategies that will minimize the potential for stigmatization, while maximizing learning and social development opportunities. A seizure action plan is specific for each student with a known seizure condition and should also include the date of last seizure, aura, length of a typical seizure for that student, frequency and how the student responds after a seizure.

Most seizures are not emergencies. The seizure action plan will help identify when a seizure is an emergency and how to respond appropriately.

Crafting the plan requires collaboration among the school nurse, student, parents/guardians, school personnel and the student's treating physicians. To ensure the student's needs are met, a seizure action plan must be student-centered and identify strategies to overcome barriers within school settings. This information may be incorporated into Individual Healthcare Plans (IHPs), 504 Plans or Individualized Education Plans (IEPs). The seizure action plan may also serve as the basis of the training curriculum for other school personnel (see Table 17).

Assessing the Student's Needs

The first steps for the school nurse in assessing a student are to gather, document and review information from the student, parents, teachers and the student's treating physician. These steps also help identify coexisting learning, behavior, mood or mobility problems that occur as a consequence of the epilepsy, the medications and any associated stigma. Completing a thorough assessment helps predict and address issues before they arise. (Please see Appendix A for forms and questionnaires that school nurses can use to collect data and structure interventions.)

TABLE 17: Key Components of a Seizure Action Plan

Assess the Student's Needs

- Gather information from parents, teachers, healthcare professionals and others.
- Observe and document seizures and circumstances (seizure precipitants and precautions)
- Observe and document responses to medications.

Customize an Action Plan

- Document treatment
- Detail first aid (care, comfort, emergency)
- Detail strategies for managing seizures in special circumstances.
- Document strategies and accommodations to address learning, behavioral and psychosocial issues.
-

Teach and Tailor Interventions as Needed

- Teach all relevant school personnel and students (first aid, relevant accommodations, and myths.)

Medication Documentation and Administration

The school nurse or her delegate is responsible for documenting and administering medications. Fortunately, many of the newer seizure medications can be given once or twice a day, often eliminating the need for the student to take medications during school hours. However, detailing daily AED administration procedures is still a critical function of the seizure action plan for most students with seizures (see Table 17).

TABLE 18: Medication Documentation and Administration

The following medication information should be documented in the seizure action plan:

- Brand and generic name of the antiepileptic medication
- Dosage
- Dosing regimen
- Special instructions for administration
- How to monitor responses
- Potential side effects
- Possible drug interactions
- What to do if a dose is missed
- What to do if side effects are observed

Documenting Seizures and Circumstances

Managing seizures in school settings begins with observing, recognizing and documenting seizure behaviors that are occurring. This can be challenging, because many symptoms are subtle, difficult to detect or occur without warning. Many are easily confused with other behavioral symptoms or problems. For example, brief staring episodes during absence seizures are often mistaken for attention disorder or daydreaming. Confusion, headaches, dizziness and language problems can also be symptoms of seizures that are often overlooked in school-aged children. (Please see page 21 for more details on conditions that might mimic epilepsy.)

Parents of students already diagnosed with epilepsy can help school personnel to observe and recognize possible seizure behaviors. They should provide the name and seizure type, detailed descriptions of the student's typical seizure(s), the date of the last seizure, the aura, usual length of a seizure, frequency and how the student responds after the seizure and first-aid measures used at home.

Non-medical school personnel need to be enlisted to help observe the student and gather information and insights into his/her seizure behaviors. Information should be gathered about what happens before, during and after the seizure.

The Epilepsy Foundation of Virginia can provide free training for school personnel on managing seizures and providing first aid. Call 434-924-8669 to schedule training.

TABLE 19: DOCUMENTING SEIZURE ACTIVITY

When a seizure is suspected, school personnel should:

- **Observe in detail the student's behaviors**
- **Time the length of the seizure**
- **Keep a written log of seizure descriptions in a communication book or flow sheet that can be shared with the school nurse, parents/guardians and healthcare team**
- **Recall and document what behaviors occurred before the seizure**
- **Determine if there is a pattern in the timing of the seizure and/or if there are triggers involved**

Behaviors Before a Seizure

Children may experience changes in mood hours - or even days - before a seizure event. Some children have a warning, or aura, such as a strange feeling, tingling, numbness or nausea that immediately precedes a seizure. Children may also report strange tastes or smells, ringing in the ears, vision changes or headache. Sudden changes in emotion (fear, anxiety, unpleasant feelings) should also be documented.

Is There a Pattern?

School personnel who observe a seizure should be instructed to look for a pattern in the timing of the seizure and possible seizure triggers. They should first note the date and time of day of each seizure. Noting and reporting all possible environmental triggers is helpful and serves to inform the seizure action planning process.

For a small number of students a specific seizure trigger may be identified. Flashing lights (photosensitivity) and hyperventilation are two documented seizure triggers.

Many other factors not considered to be true seizure triggers are also known to increase the likelihood of ~ seizure in students with epilepsy. These include the following:

- Missed or late medication (#1 reason)
- Overheating/overexertion
- Stress/anxiety
- Lack of sleep or fatigue
- Poor diet/missed meals
- Hormonal changes, particularly in adolescent girls
- Alcohol use
- Drug interactions
- Fever/colds/infection
- Low blood sugar
- Hyperventilation

A school safety evaluation should be done to assess possible seizure triggers. During this assessment it is important to ask parents about seizure triggers in their child. The assessment may reveal that activities in which the child engages may need to be modified to reduce exposure to situations that trigger seizures. For example, flashing lights in movies, television and computers may be seizure triggers. In many cases, however, there is no identifiable trigger or precipitating factor.

What Happens to the Student During a Seizure?

Teachers and others should observe and make note of the following:

- Awareness/consciousness - Is the student confused, disoriented, not responsive, unconscious?
- Speech - Is the student able to talk clearly?
Does he/she respond with only a few words or sounds? Is his/her speech nonsensical or incoherent? Is there speech arrest?
- Facial expression - Is the student staring blankly? Are eyes blinking in an unusual way? Is there facial droop or drooling?
- Turning - Is the student turning his/her head, eyes or body in an unusual way?
- Muscle tone - Is the student's body stiff or limp?
- Twitching - Is the student jerking, with movements of face, arms or legs?
- Automatism - Is the student aimlessly lip smacking, chewing, swallowing, picking at clothes, rubbing hands or tapping his/her feet?
- Walking, running or wandering - Is the student moving about without awareness of activity?
- Falling - Is the student walking unsteadily?
- Skin color - Is skin pale, cyanotic, sweating?
- Respiration - Are there breathing changes?
- Control of urine or bowels - Is the student incontinent?

What Was the Length of the Seizure?

Observers should accurately record the time from the beginning of the seizure to cessation of movement or altered behavior (ictal phase) and from the end of the seizure to the point in time when the student resumes normal activity (postictal phase). If the student transitions directly into postictal sleep, record the length of sleep.

Behavior After the Seizure?

Important observations after a seizure include:

- Is the student able to respond to voice or touch?
- Is the student aware of person, place or time?
- Is there memory of the seizure event?
- Can the student talk?
- Is there weakness or numbness in the arms or legs?
- What is the student's mood?
- Is he/she fatigued or sleepy?

Observing Responses to Treatment

The school nurse needs to gather detailed information about the student's AED regimen and the potential side effects of medication. He/she should also be aware whenever medication changes are being made. When appropriate, the school nurse should educate the student's teachers on behavioral or mood changes they may observe and establish a means of communicating these observations back to the school nurse, parents/guardians and the student's healthcare provider(s) as needed.

Assessing Learning and Psychosocial Issues

Children with seizures often experience problems with learning, behavior, mood and mobility as a consequence of the epilepsy or its underlying etiology. These consequences may impact the child in the school setting more than occasional seizures themselves. Recognition and treatment of these issues are critical to successful school experiences.

The school nurse's role in identifying and managing consequences of seizures will vary depending upon the severity of the student's condition. For example, the student with intractable seizures who experiences many injuries will need help from the school nurse to ensure that appropriate safety precautions are in place to prevent injury and allow him/her to participate fully in activities.

Evaluating the impact of seizures and medications on each student's learning and educational achievement will help identify learning or mood problems in students with seizures. The school nurse's role should focus on communicating concerns with the parents and healthcare professionals and on encouraging additional educational and medical evaluations. Strategies and accommodations to address learning deficits, mood issues and other identified disabilities should be included in both the student's seizure action plan and the IEP.

Customizing an Action Plan

Once an assessment and the collection of key information are complete the school nurse must write a customized seizure action plan for each student with seizures. The plan should be distributed to relevant school personnel. In addition to documenting seizure type and treatment, this plan should document basic and emergency firstaid procedures and list specific interventions and strategies that will minimize the potential for stigmatization while maximizing learning and social development opportunities.

Seizure first aid used to be thought of as protecting a student during a seizure or calling for emergency help. Seizure action plans, however, recognize that more can be done to help a student during a seizure and, in some circumstances, to intervene to help prevent, stop or lessen part or all of a seizure.

Educational Accommodations

The school nurse should work with teachers to make sure that they understand the dynamics of seizure disorders and their impact on the educational process for each student. Teachers should be encouraged to make accommodations to assist the student through easy-to-implement classroom practices. These accommodations may also need to be included in the student's IHP, IEP and/or 504 Plan.

TABLE 20: Simple Educational Accommodations

- Extra test time or untimed tests
- Verbal responses on tests
- Moving chair closer to the front of the room
- Tutoring or peer assistance
- Dimming fluorescent lights
- Allowing student to wear a hat if needed to dim lights
- Limit computer time

A customized seizure action plan should clearly answer the following questions:

- What medication and treatment protocol have been prescribed by the student's healthcare team?
- What is the most appropriate approach to basic first aid? This is specific to seizure type and whether consciousness is impacted.
- What is the most appropriate way to offer care and comfort after a seizure episode?
- What is normal for this student? Does the student experience an aura? Can the student usually resume normal activities within a reasonable time? If yes, the plan should define reasonable time and document who should be contacted.
- What constitutes a seizure emergency for this student? Clarify guidelines in concert with parents and the student's healthcare team.
- What is the appropriate response to an emergency? Call 911? Give PRN medications? Activate Vagus Nerve Stimulator?
- What are known seizure triggers and what precautions can be taken for avoiding them?
- What are recommended training protocols for school personnel and students?
- What aspects of the student's disorder may require referrals? For example, does the student show signs of serious depression?
- What learning, psychosocial and behavioral deficits have been identified?
- What strategies and accommodations have been implemented to support the student?

Special Circumstances and Precautions

Students with seizures should be allowed to participate in field trips, recesses and after-school activities. To ensure full participation, the school district should provide a training opportunity to all relevant staff so they can respond appropriately to seizures. The school nurse should incorporate procedures and strategies to address special needs identified by the student and his/her parents during the assessment. For example, when students with frequent seizures, a tendency to seizure clusters or long seizures participate in a field trip or afterschool activity, the school nurse may need to work creatively with the student and his/her parents to provide training for teachers and chaperones in seizure recognition and first aid so they can respond

appropriately until additional help is available. Other circumstances that may require special precautions include:

- Students involved in school sports, especially swimming and contact sports
- Students in wheelchairs
- Students who have specific, identified seizure triggers

Teaching and Tailoring Interventions

It is essential that the school nurse teach other school personnel how to respond to seizures in school settings and help to implement any special strategies and classroom or extracurricular accommodations. It is advisable to conduct a thorough assessment of the needs of the student with seizures, as well as those of teachers and other school personnel prior to implementing a training program. This will help ensure that training is focused on the most important issues. For example, teachers and other school personnel who have a student who uses a Vagus Nerve Stimulator and is going through medication changes, or has a tendency to have seizure clusters, may require special training. In addition, teachers and other school personnel who express significant fears and concerns about having a student with seizures in their classrooms may need additional training or counseling assistance.

Training protocols for school personnel must focus on dispelling myths and fears and teaching basic first aid. The training should emphasize that, while most students will not have seizures at school, they may still experience social, emotional and cognitive problems that interfere with school performance. (For more information on training school personnel, please see Section 6 of this guide.)

Interventions, whether they include training, specific accommodations or ongoing counseling, must be tailored to address the special circumstances of the student. These should be monitored and adjusted regularly to allow for medication changes and natural hormonal and growth-related shifts that can impact seizure control. Over time, additional interventions and precautions may be needed, as the original guidelines for managing each student's situation change.

This type of proactive, caring approach helps students with epilepsy to grow up to be successful, well-adjusted individuals.

Supportive Counseling and Referrals

Students with epilepsy and their families may need supportive counseling and referrals. A thorough assessment of psychosocial, learning and behavioral risk factors will help determine the best course of action. Understanding where the student is on the spectrum of severity will also help predict the likelihood of the student's developing a truly disabling condition.

The school nurse, in collaboration with a school counselor, social worker and/or school psychologist may choose to address basic behavioral, social and learning issues directly through supportive and educational counseling with the student and his/her parents. Some issues that can be addressed include:

- Medication compliance
- Dating, driving and disclosure
- Parental over-protectiveness
- Minor memory and learning impairments
- Socialization issues (anger, withdrawal, embarrassment)
- Career choices and possible work limitations

When to Refer to a Specialist


Students with more significant and severe consequences, who are not currently under the care of a specialist, may require additional referrals to medical specialists or mental health professionals. Referrals should be considered for the following:

- Uncontrolled seizures and status epilepticus
- Significant mood and anxiety disturbances
- Major memory and learning deficits
- Sudden change in seizure type
- Signs of medication toxicity or allergy

Section 6

TRAINING TEACHERS AND OTHER SCHOOL PERSONNEL

*“There is a real
need to share facts,
dispel myths and
reduce fears about
seizures.”*



Section 6

TRAINING TEACHERS AND OTHER SCHOOL PERSONNEL

The attitude and full cooperation of teachers and other school personnel are key factors in promoting positive social and educational outcomes for the student with epilepsy. Appropriate training optimizes seizure management strategies and helps ensure the full integration of the student with epilepsy into all school activities while minimizing further stigmatization.

General Curriculum for Training Others in the School Setting

Since seizures only last a few seconds to a few minutes, the seizure is often over by the time the school nurse arrives. Therefore it is important that all of those who may be in a position to assist the student with epilepsy receive information about seizure recognition, seizure first aid and the impact of seizures and epilepsy on learning and psychosocial development. This includes, but is not limited to, teachers, secretaries, athletic coaches, bus drivers, foodservice workers, administration personnel, counselors, volunteer parents and classmates.

The school nurse, the student's physician or an Epilepsy Foundation staff member or volunteer can conduct seizure recognition and first-aid training for teachers and students. A variety of training aids are available from the Epilepsy Foundation to assist with this training. These include, but are not limited to, the following:

- *Seizure Training for School Personnel:: A Presentation Guide and Toolkit*
- *Be Seizure Smart! - Teacher Information Kit*
- *Seizures and You: Take Charge of the Facts An epilepsy awareness program for teens*
- Brochures, videos, books and pamphlets targeting teens, parents, teachers and siblings
- Fact sheets on various subjects related to epilepsy and seizures, including specific treatments and classroom accommodation issues (visit www.epilepsyfoundation.org/schoolnurse to access these fact sheets)

Training Objectives

- After training, school personnel should be able to:
- Recognize seizures
- Provide appropriate first aid
- Recognize when a seizure is a medical emergency
- Provide appropriate social and academic support

Considerations in Curriculum Development

Seizure recognition and first aid should be provided for all school personnel who interact with students with seizures. Training should include at least the following information:

- What is a seizure? What is epilepsy?
- Who has epilepsy? (incidence and prevalence)
- What do seizures look like?
- What are common myths about epilepsy?
- How might seizures and epilepsy impact learning?

- When is a seizure an emergency?
- What causes seizures?
- What are common seizure triggers?
- What can be done to prevent stigma?
- What is a seizure action plan

The following optional topics might be included if relevant:

- Use of "rescue" medication including Diastat® AcuDial™
- Use of the Vagus Nerve Stimulator magnet
- Use of the ketogenic diet
- Seizures in a wheelchair
- Seizures in the water
- Seizures on a bus
- Seizures in a non-classroom area

Dispelling Myths and Reducing Fears

All school personnel should be taught the following information to dispel myths and reduce their fear of students with seizures:

- Most seizures are not medical emergencies.
- Children are usually not aware they are having a seizure and will not remember what happened.
- Epilepsy is not contagious.
- Epilepsy is not a form of mental illness.
- It is extremely unlikely that a child will die or have brain damage during a seizure.
- It is very rare for a child to become violent during a seizure.
- It is not possible to swallow one's tongue during a seizure or at any other time.

Teaching About the Impact of Epilepsy on Learning

Input from school nurses can help explain the need to modify the educational plan for some students with epilepsy. Teachers should be aware of the following information:

- Seizures often cause short-term memory problems. After a seizure event, coursework must be re-taught.
- Seizure activity in the brain, without obvious physical symptoms, can still affect learning.
- Learning will occur more easily on some days than on others.
- Medication may cause drowsiness, inattention, concentration difficulties and behavior changes.
- Children with epilepsy are more likely to suffer from depression and low self-esteem than children with diabetes or asthma.
- Children are not always aware of their seizures.
- School difficulties are not always epilepsy related.
- Each student's seizure action plan, IHP, 504 Plan and/or IEP should address these issues.

Managing Seizures in the Classroom

Teachers play a special role in setting the tone when a student experiences a seizure. If the teacher is calm and matter-of-fact and it's clear that he/she knows what to do, the student's classmates are reassured by both actions and words. The teacher should also note key information about the seizure so these details can be shared with the school nurse, the student's parents and the student's treating physician.

If the student with epilepsy and his/her parents agree to share information about the seizure with other students, then it is important that the class be able to discuss the event afterward and have a clear explanation of what happened. The following list provides key messages a nurse or teacher should convey to students who have witnessed a seizure in the classroom:

- What happened is called a seizure.
- Having a seizure is part of a health condition called epilepsy.
- No one knows exactly what causes epilepsy in some children. Fortunately, most will outgrow it.
- The seizure happened because, for just a very short time, the electrical impulses in the student's brain did not work properly and sent mixed messages to the rest of the body. Now that the seizure is over, his/her brain is working properly again.
- Epilepsy is not a disease and it cannot be caught from or given to other students.
- Epilepsy is not a mental illness or a psychological problem.
- Children who have this condition take medicine to prevent seizures, but sometimes seizures occur anyway (Even taking cough medicine does not always stop a cough).
- Seizures stop by themselves, but it is important to know first-aid steps that will keep the student safe while the seizure is happening.

Managing Seizures Under Special Circumstances

On the School Bus

The school district should allow time for the school nurse to provide basic training to all school personnel, including school bus drivers, so that they are equipped to deal with a student who experiences a seizure.

A seizure occurring on a school bus can be challenging for the school bus driver, who must safely navigate in traffic while directing students to provide first aid. During a generalized tonic-clonic seizure, the bus driver should attempt to pull over and stop the bus. The student should be placed on one side across a seating section with his/her head toward the aisle and facing away from the seat back. Alternatively, the student can be placed on one side in the aisle. Basic first-aid measures should be provided until the seizure abates and the student regains full awareness and consciousness.

Once the seizure has passed the bus driver may return to driving the bus. The bus driver's decision to return to school or proceed to the final destination should be based on school policy and the seizure action plan.

For a Student in a Wheelchair

A student in a wheelchair should remain in the chair if at all possible. First-aid responders should secure the wheelchair to prevent it from moving and gently support the student's head and body to prevent injury. Fasten the student's seatbelt (loosely) to prevent the student from falling from the wheelchair. Allow for unobstructed movement and breathing and for secretions to flow out of the mouth. If possible, pad the

wheelchair to prevent injury to the student's limbs. Follow the relevant first-aid protocol. The student's seizure action plan should be reviewed for specific first aid and emergency treatment procedures.

When Seizures Occur in Water


Remove the student from the water immediately, supporting the head so that both the mouth and nose are always above the water. If the student is not breathing, begin rescue breathing. Always transport the student to the emergency room even if he or she appears fully recovered.

In Non-Classroom Areas of the School

For each area of the school (hallway, cafeteria, gymnasium, playground, playing field, auditorium, restroom), there should be a trained first responder and a backup first responder listed in the seizure action plan procedures for seizure response. Responsible students as well as adults can be trained as first responders.

Section 7 EPILEPSY RESOURCES

“School nurses and educators should act as role models for other school personnel and students, helping them see that seizures are a medical problem that can be treated easily.”



Section 7

EPILEPSY RESOURCES

The resources listed in this section will be of value to the school nurse dealing with students with seizures. For more details and an up-to-date list of resources, school nurses can visit a special section on the Epilepsy Foundation Web site at the following:

www.epilepsyfoundation.org/schoolnurse

Community Resources

There are many Epilepsy Foundation affiliates serving local communities throughout the United States. Many books, kits, manuals, pamphlets, posters and videos targeting health professionals, parents, children, teens and other populations are available through these offices. To locate the affiliate nearest you, please call the national office of the Epilepsy Foundation at

(800)332-1000 or visit www.epilepsyfoundation.org/aboutus/AffiliateLookup.cfm

The Epilepsy Foundation Office of Virginia office is located at 500 Ray C. Hunt Drive, Charlottesville, Virginia 22903. The mailing address is P.O. Box 800659, UV A Health Center, Charlottesville, Virginia 22908.

Telephone is 434-924-8669.

Website is <http://www.efva.org>

Five regional offices and director names, addresses and contact information can be found at <http://www.epilepsyfoundation.org/local/virginia/>

Suzanne Bischoff is the Executive Director and can be emailed at srb3m@virginia.edu

Internet Resources

Below is a short list of valuable Internet resources. For a more exhaustive and updated list please visit www.epilepsyfoundation.org/schoolnurse.

The Epilepsy Foundation

www.epilepsyfoundation.org

The Epilepsy Foundation is the national voluntary agency solely dedicated to the welfare of the 3 million people with epilepsy in the U.S. and their families. The organization works to ensure that people with seizures are able to participate in all life experiences; and to prevent, control and cure epilepsy through research, education, advocacy and services.

The Epilepsy Foundation Virginia

www.efva.org

This is our state affiliate solely dedicated to the welfare of the 800,00 people with epilepsy in Virginia and their families. The organization works to ensure that people with seizures are able to participate in all life experiences; and to prevent, control and cure epilepsy through research, education, advocacy and services.

American Epilepsy Society

www.aesnet.org

The American Epilepsy Society promotes research and education for professionals dedicated to the prevention, treatment and cure of epilepsy.

Epilepsy.com

www.epilepsy.com

Epilepsy.com is an online resource provided by The Epilepsy Project. Its mission is to inform and empower two groups of patients and their families: those facing newly diagnosed epilepsy and those struggling with epilepsy that has resisted the usual treatments.

American Academy of Neurology

www.aan.com

The American Academy of Neurology provides valuable resources for medical specialists worldwide who are committed to improving the care of patients with neurological diseases.

American Clinical Neurophysiology Society

www.acns.org

The ACNS is a professional association dedicated to fostering excellence in clinical neurophysiology and furthering the understanding of central nervous system function in health and disease through education, research and the provision of a forum for discussion and interaction

American Neurological Association

www.aneuroa.org

The American Neurological Association is a professional society of academic neurologists and neuroscientists devoted to advancing the goals of academic neurology, to training and educating neurologists and other physicians in the neurological sciences and to expanding both our understanding of diseases of the nervous system and our ability to treat them.

Care Connection

www.vahealth.org/specialchildren/directory/health.asp

This directory, created for parents, children, health providers and community organizations, allows you to search a list of local and Virginia resources for contact information on various medical conditions, advocacy and support groups, insurance, community support organizations and public and private agencies providing services for children with special health care needs.

Centers for Disease Control and Prevention Epilepsy Web Site

www.cdc.gov/epilepsy/index.htm

The national Centers for Disease Control and Prevention is recognized as the lead federal agency for protecting the health and safety of people at home and abroad, providing credible information to enhance health decisions and promoting health through strong partnerships. Its epilepsy web site details current partnerships and health promotion activities being supported by the agency's epilepsy program.

The Charlie Foundation

www.charlifoundation.org

The Charlie Foundation was established In 1994 to raise awareness about the ketogenic diet as a treatment for childhood epilepsy. It facilitates basic and clinical research, educates professionals and informs families about the current status of the ketogenic diet.

The International Bureau for Epilepsy

www.ibe-epilepsy.org

The International Bureau for Epilepsy was established in 1961 as an organization of laypersons and professionals interested in the medical and non-medical aspects of epilepsy. The IBE addresses such social problems as education, employment, insurance, driver's license restrictions and public awareness.

The International League Against Epilepsy

www.ilae-epilepsy.org

The International League Against Epilepsy works to advance and disseminate knowledge concerning epilepsy. Membership is comprised of national professional organizations and individuals involved in research and those interested in the exchange of scientific information concerning epilepsy.

The National Institute of Neurological Disorders and Stroke

www.ninds.nih.gov

The National Institute of Neurological Disorders and Stroke supports and conducts research on brain and nervous system disorders. By supporting and conducting neurological research, the NINDS seeks better understanding, diagnosis, treatment and prevention of these disorders.

Neuroscience for Kids

Faculty.washington.edu/chudler/neurok.html

Neuroscience for Kids was created for all students and teachers who would like to learn about the nervous system.

PACE (Parents Against Childhood Epilepsy, Inc.)

www.paceusa.org

PACE was founded in 1996 by a group of parents in response to their experiences in caring for the medical, physical, social, educational developmental and emotional needs of their children who have epilepsy and severe seizure disorders.

Fact Sheets on Epilepsy

The Epilepsy Foundation has fact sheets on a variety of epilepsy-related topics including seizure-specific information, treatment, education, insurance and employment. A complete list of fact sheets can be accessed from the school nurse web page at www.epilepsyfoundation.org/schoolnurse.

Many of the fact sheets can be downloaded for your immediate use. Please check the website regularly as this list is periodically updated.

- Below is a partial list of topics covered:
- Alcohol and Drug Abuse
- Blood-Level Monitoring of Antiepileptic Medication
- Diagnosing Epilepsy and Seizure Disorders
- Effects of Epilepsy on Learning and Memory
- Financial Aid: Scholarships, Grants and Fellowships
- IDEA: Elementary and Secondary Education
- Insurance Issues in Epilepsy and Seizure Disorders
- Nonepileptic Seizures
- Patterns of Seizure Frequency
- Photosensitivity and Seizures
- Prognosis in Epilepsy and Seizure Disorders
- Remission vs. Cure for Epilepsy
- Seizure Classifications: Absence Seizures
- Seizure Classifications: Complex Partial Seizures
- Seizure Classifications: Tonic-Clonic Seizures

Driving and Epilepsy

Every state regulates driver's license eligibility for those with certain medical conditions. The most common requirement for those with epilepsy is that they be seizure-free for a specific period of time and submit a physician's evaluation of their ability to drive safely.

In Virginia, the law requires the driver to be seizure free for a period of six months, with some exceptions. Persons with nocturnal seizures may drive, but are subject to review every three months. A person who has been seizure free for two years may drive without any restrictions. Medical personnel are not required to report epilepsy to DMV. If the person already has a license, it will be suspended for six months from the date of the last seizure/blackout. For more information go to the Virginia DMV website: <http://www.dmv.state.va.us/>

BIBLIOGRAPHY

Bibliography

In our school nurse program, extensive use has been made of the 2009 school nurse manual from the Epilepsy Foundation in Landover, MD. We thank Jody Kakacek and others for their cooperation in this matter.

Section 1

Delorenzo RJ, Towne AR, Pellock JM, Ko D. Status epilepticus in children, adults, and the elderly. *Epilepsia*. 1992;33(Suppl4):S15-S25.

Devinsky O. *Epilepsy: Patient 6- Family Guide*. Philadelphia, Pa: EA. Davis Company; 2002.

Glauser T. Pediatric epilepsy syndromes. *Current Opinions in Pediatrics*. 1995;7:640-649.

Hauser A. Epidemiology of epilepsy. In: Sanntilli N, ed. *Managing Seizure Disorders*. Philadelphia, Pa.: Lippincott-Raven; 1996:7-18.

Hauser A. Epidemiology of epilepsy in children. In: Pellock J, Dodson W, Bourgeois B, eds. *Pediatric Epilepsy: Diagnosis and Therapy* New York: Demos, 2001:81-96.

McDermott S, Moran R. Prevalence of Epilepsy in Adults with Mental Retardation and Related Disabilities in Primary Care. *American Journal on Mental Retardation*. January; 2005 - Vol. 10, No. 1 :48-56.

Palencia R. Prevalence and incidence of epilepsy in childhood. *Revista de Neurologia* 2000; Suppl 1: SI-4.

Prensky A. An approach to the child with paroxysmal phenomena with emphasis on nonepileptic disorders. In: Pellock J, Dodson W, Bourgeois B, eds. *Pediatric Epilepsy: Diagnosis and Therapy*. New York: Demos, 2001:97-115.

Smith EA, VanHauten R. A comparison of the characteristics of self-stimulating behaviors in "normal" children and children with developmental delay. *Res Dev Disability*. 1996; 17:253-268.

Section 2

Drone DL, Meador KJ. Cognitive and behavioral effects of AEDs. *Epilepsy and Behavior*. Oct 2002; 3; 2: S49-S53.

Hermann Bp, Austin J. Psychosocial status of children with epilepsy and the effects of epilepsy surgery. In:

Wyllie E, ed. *The Treatment of Epilepsy Principles and Practices*. Philadelphia, Pa.: Lea and Febiger; 1993: 99:1141 -1148.

Fastenau, P.S., Shen, J., Dunn, D.W. & Austin, J.K. Academic underachievement among children with epilepsy: proportion exceeding psychometric criteria for learning disability and associated risk factors. *Journal of Learning Disability* 2008; 41(3):195-207.

Prevey M, Delancey RJ, Cramer R et al. Epilepsy Cooperative study 264 group. Complex partial and secondarily generalized seizure patients: Cognitive functioning prior to treatment with anti epileptic medicine. *Epilepsy Research*. 1998; 30:1-9.

Sutula Tp, Hagen J, Pitkanen A. Do Epileptic Seizures Damage the Brain? *Curr Opin Neurology*. 2003; 16:189-195.

Sabaz M, Cairns D, Lawson JA, et al. The health-related quality-of-life of children with refractory epilepsy: A comparison of those with and without intellectual disability. *Epilepsia*. 2001; 42: 5:621-628.

Sperling, M. The Consequences of Uncontrolled Epilepsy. *CNS Spectrums*. 2004; 9; 2:98-108.

Swartztrauber K, Dewar S and Engel J. Patient attitudes about treatment for intractable Epilepsy. *Epilepsy and Behavior*. 2003, 4: 19-25.

Wiebe S. Effectiveness and safety of epilepsy surgery. What is the evidence? *CNS Spectrums*. 2004; 9; 2:120-132.

Zelnick N, Salado L et al. Seizure control and educational outcome in childhood onset epilepsy. *J Child Neurol*. 2001; 16:820-24.

Section 4

Harbord MG, Kyrkou MR, Kay O, Coulthard KP. Use of intranasal midazolam to treat acute seizures in paediatric community settings. *J Paediatr Child Health* 2004; 40(9-10):556-8.

APPENDICES

Appendices

Appendix A:

Seizure Action Planning Forms (snapshots – to print forms to be used please see PDF files on disc)

Appendix B:

Seizure Recognition and First-Aid Chart (snapshots – to print charts to be used please see PDF files on disc)

Appendix C:

Antiepileptic Medications Chart(snapshots – to print charts to be used please see PDF files on disc)

Appendix A:

Seizure Action Planning Forms

The sample forms that follow provide guidelines for collecting data relevant to educational and seizure action planning for students with epilepsy. They can be adapted to meet individual and institutional needs. Each form provides a framework for collecting and sharing information critical to appropriate health management planning, which can occur only if the data are used collectively.

As information is gathered, transportation issues need to be considered. An assessment of school safety, including the playground and gym activities needs to be completed. The best method for communicating with parents and medical team also needs to be determined.

1. Questionnaire for Parent

- can be difficult to get parents to complete the form and may require persistence
- may require interview the parent to clarify the information
- update annually or whenever changes occur
- will assist school personnel in recognizing a student's seizure and responding appropriately

2. Seizure Observation Record

- to be completed by school personnel when reporting a seizure
- helps identify seizure triggers, patterns and precautions
- needs to be communicated with parents

3. Seizure Action Plan

- to be completed by school nurse at earliest time
- provides basic information about student's seizures and medication
- distribute to relevant school personnel should be approved by parents and physician



Please complete all questions. This information is essential for the school nurse and school staff in determining your child's special needs and providing a positive and supportive learning environment. If you have any questions about how to complete this form, please contact your child's school nurse.

Contact Information

Student's Name	School Year	Date of Birth	
School	Grade	Classroom	
Parent/Guardian	Phone	Work	Cell
Parent/Guardian Email			
Other Emergency Contact	Phone	Work	Cell
Child's Neurologist	Phone	Location	
Child's Primary Care Doctor	Phone	Location	
Significant Medical History or Conditions			

Seizure Information

1. When was your child diagnosed with seizures or epilepsy? _____

2. Seizure type(s) _____

Seizure Type	Length	Frequency	Description

3. What might trigger a seizure in your child? _____

4. Are there any warnings and/or behavior changes before the seizure occurs? ☐ YES ☐ NO
If YES, please explain: _____

5. When was your child's last seizure? _____

6. Has there been any recent change in your child's seizure patterns? ☐ YES ☐ NO
If YES, please explain: _____

7. How does your child react after a seizure is over? _____

8. How do other illnesses affect your child's seizure control? _____

Basic First Aid: Care & Comfort

9. What basic first aid procedures should be taken when your child has a seizure in school?

10. Will your child need to leave the classroom after a seizure? ☐ YES ☐ NO
If YES, what process would you recommend for returning your child to classroom: _____

Basic Seizure First Aid

- Stay calm & track time
 - Keep child safe
 - Do not restrain
 - Do not put anything in mouth
 - Stay with child until fully conscious
 - Record seizure in log
- For tonic-clonic seizure:**
- Protect head
 - Keep airway open/watch breathing
 - Turn child on side

Seizure Emergencies

11. Please describe what constitutes an emergency for your child? (Answer may require consultation with treating physician and school nurse.)

12. Has child ever been hospitalized for continuous seizures? ☐ YES ☐ NO

If YES, please explain:

A seizure is generally considered an emergency when:

- Convulsive (tonic-clonic) seizure lasts longer than 5 minutes
- Student has repeated seizures without regaining consciousness
- Student is injured or has diabetes
- Student has a first-time seizure
- Student has breathing difficulties
- Student has a seizure in water

Seizure Medication and Treatment Information

13. What medication(s) does your child take?

Medication	Date Started	Dosage	Frequency and Time of Day Taken	Possible Side Effects

14. What emergency/rescue medications are prescribed for your child?

Medication	Dosage	Administration Instructions (timing* & method**)	What to Do After Administration

* After 2nd or 3rd seizure, for cluster of seizure, etc.

** Orally, under tongue, rectally, etc.

15. What medication(s) will your child need to take during school hours?

16. Should any of these medications be administered in a special way? ☐ YES ☐ NO

If YES, please explain:

17. Should any particular reaction be watched for? ☐ YES ☐ NO

If YES, please explain:

18. What should be done when your child misses a dose?

19. Should the school have backup medication available to give your child for missed dose? ☐ YES ☐ NO

20. Do you wish to be called before backup medication is given for a missed dose? ☐ YES ☐ NO

21. Does your child have a Vagus Nerve Stimulator? ☐ YES ☐ NO

If YES, please describe instructions for appropriate magnet use:

Special Considerations & Precautions

22. Check all that apply and describe any consideration or precautions that should be taken:

- | | |
|---|--|
| <input type="checkbox"/> General health | <input type="checkbox"/> Physical education (gym/sports) |
| <input type="checkbox"/> Physical functioning | <input type="checkbox"/> Recess |
| <input type="checkbox"/> Learning | <input type="checkbox"/> Field trips |
| <input type="checkbox"/> Behavior | <input type="checkbox"/> Bus transportation |
| <input type="checkbox"/> Mood/coping | <input type="checkbox"/> Other |

General Communication Issues

23. What is the best way for us to communicate with you about your child's seizure(s)?

24. Can this information be shared with classroom teacher(s) and other appropriate school personnel? ☐ YES ☐ NO

Dates

Updated

Parent/Guardian Signature _____ Date _____

DPC775



Seizure Action Plan

Effective Date _____

This student is being treated for a seizure disorder. The information below should assist you if a seizure occurs during school hours.

Student's Name _____	Date of Birth _____
Parent/Guardian _____	Phone _____ Cell _____
Other Emergency Contact _____	Phone _____ Cell _____
Treating Physician _____	Phone _____
Significant Medical History _____	

Seizure Information

Seizure Type	Length	Frequency	Description

Seizure triggers or warning signs: _____

Student's response after a seizure: _____

Basic First Aid: Care & Comfort

Please describe basic first aid procedures: _____

Does student need to leave the classroom after a seizure? ☐ Yes ☐ No

If YES, describe process for returning student to classroom: _____

Emergency Response

A "seizure emergency" for this student is defined as: _____

Seizure Emergency Protocol

(Check all that apply and clarify below)

- ☐ Contact school nurse at _____
- ☐ Call 911 for transport to _____
- ☐ Notify parent or emergency contact
- ☐ Administer emergency medications as indicated below
- ☐ Notify doctor
- ☐ Other _____

Basic Seizure First Aid

- Stay calm & track time
- Keep child safe
- Do not restrain
- Do not put anything in mouth
- Stay with child until fully conscious
- Record seizure in log

For tonic-clonic seizure:

- Protect head
- Keep airway open/watch breathing
- Turn child on side

A seizure is generally considered an emergency when:

- Convulsive (tonic-clonic) seizure lasts longer than 5 minutes
- Student has repeated seizures without regaining consciousness
- Student is injured or has diabetes
- Student has a first-time seizure
- Student has breathing difficulties
- Student has a seizure in water

Treatment Protocol During School Hours (include daily and emergency medications)

Emerg. Med. ✓	Medication	Dosage & Time of Day Given	Common Side Effects & Special Instructions

Does student have a Vagus Nerve Stimulator? ☐ Yes ☐ No If YES, describe magnet use: _____

Special Considerations and Precautions (regarding school activities, sports, trips, etc.)

Describe any special considerations or precautions: _____

Physician Signature _____ Date _____

Parent/Guardian Signature _____ Date _____

DPC772



Seizure Observation Record

Student Name:			
Date & Time			
Seizure Length			
Pre-Seizure Observation (Briefly list behaviors, triggering events, activities)			
Conscious (yes/no/altered)			
Injuries? (briefly describe)			
Muscle Tone/Body Movements	Rigid/clenching		
	Limp		
	Fall down		
	Rocking		
	Wandering around		
	Whole body jerking		
Extremity Movements	(R) arm jerking		
	(L) arm jerking		
	(R) leg jerking		
	(L) leg jerking		
	Random Movement		
Color	Bluish		
	Pale		
	Flushed		
Eyes	Pupils dilated		
	Turned (R or L)		
	Rolls up		
	Staring or blinking (clarify)		
	Closed		
Mouth	Salivating		
	Chewing		
	Lip smacking		
Verbal Sounds (gagging, talking, throat clearing, etc.)			
Breathing (normal, labored, stopped, noisy, etc.)			
Incontinent (urine or feces)			
Post-Seizure Observation	Confused		
	Sleepy/tired		
	Headache		
	Speech slurring		
	Other		
Length to Orientation			
Parents Notified? (time of call)			
EMS Called? (call time & arrival time)			
Observer's Name			

Please put additional notes on back as necessary.

Appendix B:

Seizure Recognition and First-Aid Charts

- **Seizure First Aid Non-Convulsive – English**
- **Seizure First Aid Non-Convulsive- Spanish**
- **Seizure First Aid Convulsive- English**
- **Seizure First Aid Non-Convulsive- Spanish**

First Aid for Seizures

(Complex partial, psychomotor, temporal lobe)

1. Recognize common symptoms



2. Follow first-aid steps



People who've had this type of seizure should be fully conscious and aware before being left on their own. Make sure they know the date, where they are, and where

they're going next. Confusion may last longer than the seizure itself and may be hazardous. If full awareness does not return, call for medical assistance.



**EPILEPSY
FOUNDATION®**

Not another moment lost to seizures™

1-800-332-1000 • www.epilepsyfoundation.org

EFA 341

© 2005 Epilepsy Foundation of America, Inc.

This publication was made possible by a grant from the Centers for Disease Control and Prevention (grant number 5U58DP00606-03) and its contents are solely the responsibility of its authors and do not necessarily represent the official views of the CDC.

Primeros auxilios para convulsiones

(Crisis parciales complejas, psicomotrices, de lóbulo temporal)

1. Reconocer los síntomas comunes



2. Seguir los pasos de primeros auxilios



Antes de que se les pueda dejar solas, las personas que han tenido este tipo de convulsión deben estar completamente conscientes y haber recobrado el conocimiento. Asegúrese de que sepan la fecha,

dónde están y adónde se dirigen. La confusión puede ser peligrosa porque a veces dura más que la convulsión propiamente. Si la persona no recupera el conocimiento por completo, solicite asistencia médica.


EPILEPSY FOUNDATION®
Not another moment lost to seizures™
 1-866-748-8008 • www.fundacionparalaepilepsia.org

EFA 341

La producción de este folleto ha sido posible por el auxilio #5U58CP00606-03 de los Centros para el Control y Prevención de Enfermedades (CDC por sus siglas en inglés) y su contenido es únicamente responsabilidad de los autores y no representa necesariamente el punto de vista del CDC.
 © 2006 Epilepsy Foundation of America, Inc.

First Aid for Seizures

(Convulsive, generalized tonic-clonic, grand mal)



Most seizures in people with epilepsy are not medical emergencies. They end after a minute or two without harm and usually do not require a trip to the emergency room.

But sometimes there are good reasons to call for emergency help. A seizure in someone who does not have epilepsy could be a sign of a serious illness.

Other reasons to call an ambulance include:

- A seizure that lasts more than 5 minutes
- No "epilepsy" or "seizure disorder" identification
- Slow recovery, a second seizure, or difficulty breathing afterwards
- Pregnancy or other medical diagnosis
- Any signs of injury or sickness

EPILEPSY FOUNDATION®
Not another moment lost to seizures™
 1-800-332-1000 • www.epilepsyfoundation.org

EFA 342

© 2008 Epilepsy Foundation of America, Inc.

This publication was made possible by a grant from the Centers for Disease Control and Prevention (grant number 5U58DP00606-03) and its contents are solely the responsibility of its authors and do not necessarily represent the official views of the CDC.

Primeros auxilios para convulsiones (Convulsiones tónicoclónicas generalizadas)



La mayoría de las convulsiones que acontecen a las personas con epilepsia no constituyen emergencias médicas. Generalmente duran sólo uno o dos minutos sin causar daños y no suelen requerir una visita a la sala de emergencias.

Sin embargo, a veces existen buenas razones para solicitar ayuda de emergencia. En personas que no tienen epilepsia, una convulsión podría ser señal de enfermedad grave.

Otras razones para llamar a una ambulancia:

- La convulsión dura más de 5 minutos
- No se localiza una identificación que asiente: epilepsia o trastorno convulsivo
- Recuperación lenta, una segunda convulsión o dificultad para respirar después de la crisis
- Embarazo u otro diagnóstico médico
- Cualquier señal de lesión o enfermedad

 **EPILEPSY
FOUNDATION®**
Not another moment lost to seizures™
1-866-748-8008 • www.fundacionparalaepilepsia.org

EFA 342

La producción de este folleto ha sido posible por el auxilio #5U58DP00006-03 de los Centros para el Control y Prevención de Enfermedades (CDC por sus siglas en inglés) y su contenido es únicamente responsabilidad de los autores y no representa necesariamente el punto de vista del CDC.
© 2009 Epilepsy Foundation of America, Inc.

Appendix C:

Antiepileptic Medications Chart

Medicines for Epilepsy

This chart is designed to help people with epilepsy (seizure disorders) become more familiar with the medications they are taking. It is not designed for use by health or other professionals to identify drugs. Other drugs not listed here may also be prescribed to prevent seizures.

The pictures are of brand name drugs as of the printing date. However, drugs may change in appearance from time to time. Generic versions of these drugs (listed in italics, under the brand name) will look different from the brand name drugs.

The list of side effects is not complete because of space limitations. For a complete list, consult your doctor, nurse,

FORM (of brand name product)	BRAND NAME (generic name - images not shown)	AVERAGE ADULT DAILY DOSE <small>(usual dosages given with epilepsy may be higher or lower than the dose listed)</small>	SOME SIDE EFFECTS <small>(Not all individuals experience side effects. The potential for adverse side effects includes any reactions.)</small>
	ATTIVAN® (clonazepam)	1 mg - 10 mg	Drowsiness, dizziness, fatigue, poor coordination, unsteadiness, behavior changes
	CARBATROL® (extended release carbamazepine)	600 mg - 1200 mg	Dizziness, drowsiness, blurred or double vision, nausea, skin rashes, abnormal blood counts (rare)
	DEPAKENE® (valproate)	1750 mg - 3000 mg	Upset stomach, abnormal bleeding time, liver toxicity, hair loss, weight gain, tremor
	DEPAKOTE® (divalproex sodium)	1750 mg - 3000 mg	Upset stomach, abnormal bleeding time, liver toxicity, hair loss, weight gain, tremor
	DEPAKOTE® ER (extended release divalproex sodium)	2000 mg - 3500 mg	Upset stomach, abnormal bleeding time, liver toxicity, hair loss, weight gain, tremor
	DIAMOX® SEQUEL® (extended release acetazolamide)	250 mg - 1000 mg	Appetite loss, frequent urination, drowsiness, confusion, numbness of extremities, kidney stones
	DILANTIN® (phenytoin)	200 mg - 600 mg	Chills, insomnia, motor twitching, rashes, rash, gum overgrowth, halitosis, thickening of fingers
	FELBATOL® (felbamate)	1200 mg - 3600 mg	Anorexia, vomiting, insomnia, nausea, headache, liver and blood toxicity
	GABRIL® (gabapentin)	32 mg - 56 mg	Tremor, dizziness, nervousness, difficulty concentrating, sleepiness, weakness
	KEPPRA® (levetiracetam)	1000 mg - 3000 mg	Sleepiness, fatigue, poor coordination, loss of strength, dizziness
	Keppra XR® (extended release levetiracetam)	1000 mg - 3000 mg	Sleepiness, fatigue, poor coordination, loss of strength, dizziness
	KLONOPIN® (clonazepam)	1.5 mg - 20 mg	Drowsiness, dizziness, fatigue, poor coordination, unsteadiness, behavior changes
	LAMICTAL® (lamotrigine)	100 mg - 500 mg	Dizziness, headache, blurred vision, chills, drowsiness, nausea, skin rash

Medicines for Epilepsy

This chart is designed to help people with epilepsy (seizure disorders) become more familiar with the medications they are taking. It is not designed for use by health or other professionals to identify drugs. Other drugs not listed here may also be prescribed to prevent seizures.

The pictures are of brand name drugs as of the printing date. However, drugs may change in appearance from time to time. Generic versions of these drugs (listed in italics, under the brand name) will look different from the brand name drugs.

The list of side effects is not complete because of space limitations. For a complete list, consult your doctor, nurse,

FORM (of brand name product)	BRAND NAME (generic name - images not shown)	AVERAGE ADULT DAILY DOSE <small>(based on body weight and epilepsy type; higher or lower dose may be used)</small>	SOME SIDE EFFECTS <small>(Not all individuals experience side effects. The patient should consult their doctor if they experience any symptoms.)</small>
	ATTIVAN® (clonazepam)	1 mg - 10 mg	Drowsiness, sleepiness, fatigue, poor coordination, unsteadiness, behavior changes
	CARBATROL® (extended release carbamazepine)	600 mg - 1200 mg	Dizziness, drowsiness, blurred or double vision, nausea, skin rashes, abnormal blood counts (rare)
	DEPAKENE® (sodium valproate)	1750 mg - 3000 mg	Upset stomach, altered bleeding time, liver toxicity, hair loss, weight gain, tremor
	DEPAKOTE® (divalproex sodium)	1750 mg - 3000 mg	Upset stomach, altered bleeding time, liver toxicity, hair loss, weight gain, tremor
	DEPAKOTE® ER (extended release divalproex sodium)	2000 mg - 3500 mg	Upset stomach, altered bleeding time, liver toxicity, hair loss, weight gain, tremor
	DIAMOX® SEQUEL® (extended release acetazolamide)	250 mg - 1000 mg	Appetite loss, frequent urination, drowsiness, confusion, numbness of extremities, kidney stones
	DILANTIN® (phenytoin)	200 mg - 400 mg	Chills, fever, insomnia, muscle twitching, nausea, rash, gum overgrowth, hiccups, thickening of fingers
	FELBATOL® (felbamate)	1200 mg - 3600 mg	Anorexia, vomiting, insomnia, nausea, headache, liver and blood toxicity
	GABITRIL® (gabapentin)	32 mg - 56 mg	Tiredness, dizziness, nervousness, difficulty concentrating, sleepiness, weakness
	KEPPRA® (levetiracetam)	1000 mg - 3000 mg	Sleepiness, fatigue, poor coordination, loss of strength, dizziness
	Keppra XR® (extended release levetiracetam)	1000 mg - 3000 mg	Sleepiness, fatigue, poor coordination, loss of strength, dizziness
	KLONOPIN® (clonazepam)	1.5 mg - 20 mg	Drowsiness, sleepiness, fatigue, poor coordination, unsteadiness, behavior changes
	LAMICTAL® (lamotrigine)	100 mg - 500 mg	Dizziness, headache, blurred vision, chills, sleepiness, nausea, skin rash




Newer Anti-Epileptic Drugs (AEDs)

Images (Generic brands may be different)	Name of Drug (Generic Name)	Range of Daily Dosage	Common Side Effects
	Banzel (Rufinamide)	400mg-3200mg	Dizziness, fatigue, difficulty concentrating, vomiting, irritability, drowsiness, appetite loss, headache
	Celontin (Methsuximide)	300mg-1.2g	Nausea, vomiting, loss of appetite, headache, drowsiness, dizziness, difficulty concentrating, confusion, blurred vision, poor coordination
image not available	Clobazam (Onfi)	20mg-40mg	Tiredness, sedation, aggression, insomnia, physical and psychological dependence
	Depacon (Valproate Sodium Injection)	10mg/kg-60mg/kg	Dizziness, headache, insomnia, sleepiness, blurred vision, vomiting, nausea, diarrhea, tremors, appetite loss
	Diasat (Diazepam Rectal Gel)	5mg-20mg	Behavior changes, sleepiness, unsteadiness, drowsiness, fatigue, poor coordination, upset stomach, insomnia, confusion, nervousness, dizziness
	Mebaral (Mephobarbital)	48mg-600mg	Poor coordination, dizziness, drowsiness, headache, nausea, vomiting, fatigue, vertigo
	Peganone (Ethotoin)	500mg-3g	Diarrhea, dizziness, nausea, vomiting
image not available	Potiga (Ezogabine) *available 2012	600mg-1200mg	Dizziness, fatigue, confusion, tremor, poor coordination, memory problems, difficulty concentrating, double vision, urinary retention
	Sabril (Vigabatrin)	2g- 4g	Vision loss, dizziness, drowsiness, nausea, vomiting, headache, insomnia, poor coordination, joint pain, memory problems

This chart is designed to help people with epilepsy (seizure disorders) become more familiar with the medication they are taking. It is not designed for use by health or other professionals to identify drugs. Other drugs not listed here may also be prescribed to prevent seizures.

The pictures are of brand name drugs as of the published date. However, drugs may change in appearance from time to time. Generic versions of these drugs will look different from the brand name drugs.

The list of side effects is not complete because of space limitation. For a complete list, consult your doctor, nurse or pharmacist.

	Tridione (Trinethaione)	900mg-2400mg	Drowsiness, upset stomach, loss of appetite, headache, irritability, fatigue
	Valium (Diazepam)	2mg-10mg	Sleepiness, fatigue, poor coordination, confusion, blurred vision, difficulty speaking
	Vimpat (Lacosamide)	200mg-400mg	Vomiting, nausea, drowsiness, tremor, dizziness, blurred vision, headache

Epilepsy Foundation of America, 2011

Index

Absence Seizures.....	2, 14, 17, 67
Accommodations	4, 53, 54
Acute Interventions	3, 32, 34
Alternative Therapies.....	4, 46
Antiepileptic Drugs	4, 39, 42
Atonic Seizures.....	2, 15
Automatisms	13, 52
Basic First Aid	3, 32
Behavior Contracts.....	29
Behavioral Concerns.....	3, 29
Benign Rolandic Epilepsy	2, 16
Cerebral Palsy.....	10, 11
Childhood Absence Epilepsy	2, 17
<i>Clonic Seizures</i>	2, 15, 67
Clusters.....	14, 17, 23, 33, 35, 41, 54, 55
Cognitive Dysfunction	27
Complex Partial Seizures.....	13, 18, 29, 33
Counseling	5, 55
Delegation	3, 37
Driving	5, 67
Drop Attack	18
Emergency Response	3, 32, 34
Epilepsy Foundation Of Virginia.....	2, 6, 7
Epileptic Syndromes.....	2, 16
Female.....	3, 23
Frontal Lobe Epilepsy	18
Frontal Lobe Epilepsy	2, 18
Generalized Seizures	2, 14
Impact Of Epilepsy	3, 5, 26, 27, 28, 60
Intractable Epilepsy	3, 21
Juvenile Myoclonic Epilepsy	2, 17
Ketogenic Diet.....	4, 43, 44
Landau-Kleffner Syndrome.....	2, 19
Learning, Impact Of Epilepsy.....	3, 4, 5, 26, 53, 60, 67, 68
Lennox-Gastaut Syndrome	2, 17
Managing Seizures	
On The School Bus, In A Wheelchair, In	
Nonclassroom Areas, In The Classroom,	
In Water	5, 61
Managing Stress	4, 46
Memory Problems	18
Menstrual Cycles.....	23
Mental Retardation.....	10, 11, 17, 18, 27
Modified Atkins Diet.....	4, 44
<i>Myoclonic Seizures</i>	2, 15
Myths.....	5, 60
Non-Epileptic Seizures.....	3, 20
Partial Seizures.....	12, 13, 16, 18, 23, 32, 43
Photosensitivity.....	51
Photosensitivity.....	67
Prn Medications	4, 41
Progressive Myoclonic Epilepsy	2, 18
Pseudoseizures	20
Psychogenic Seizures	20
Quality Of Life	8, 22, 25, 26
Reflex Epilepsy	2, 19
Resources	5, 64
Seizure Action Plans.....	4, 49
Seizure-Like Phenomena	3, 20
Self-Esteem	22, 27, 60
Service Dogs	4, 45
Side Effects	21, 25, 26, 28, 29, 36, 38, 39, 40, 41, 42, 43, 44, 53
Simple Partial Seizures.....	13
Spectrum Of Severity	29, 55
Status Epilepticus.....	3, 22, 23
Stigma.....	3, 29
Surgery	4, 39, 43
Syncope	20
Teenage	3, 30
Temporal Lobe Epilepsy	2, 18
<i>Tonic Seizures</i>	2, 15
Training Teachers	5, 59
Treatment	3, 39
Triggers.....	19, 46, 51, 52, 54, 60, 70
Vagus Nerve Stimulator (Vns)	44..... 4

Index